

Solving Multiobjective Optimization Problems with inequality constraint using an augmented Lagrangian function

Appolinaire Tougma, Kounhinir Somé *

*Département de Mathématiques, Laboratoire de Mathématiques, Informatique et Applications (L@MIA),
Université Norbert Zongo, BP 376 Koudougou, Burkina Faso*

Abstract

This paper presents a technique for addressing multiobjective optimization issues subject to inequality constraints. This technique converts the original problem into a single-objective optimization without constraints by employing an augmented Lagrangian function and an ϵ -constraint method. Specifically, the augmented Lagrangian function transforms problems with multiple objectives into a single objective function, while the ϵ -constraint method changes constrained optimization problems into unconstrained ones. We provide two propositions complete with proofs to verify the admissibility and Pareto optimality of the solutions derived. Furthermore, we conduct a comparative analysis with two established methods, NSGA-II and BoostDMS, focusing on the convergence and distribution of solutions across fifty test problems sourced from existing literature. The collective theoretical and empirical evidence suggests that our proposed method is superior for solving multiobjective optimization problems.

Keywords Multiobjective optimization, Penalty function, Augmented Lagrangian, Gradient projected, ϵ -Constraint approach

AMS 2020 subject classifications 49M37, 65K05, 90C25, 90C29

DOI: 10.19139/soic-2310-5070-1875

1. Introduction

Recently, the application of multiobjective optimization concepts has been instrumental in addressing specific challenges in various domains, including physics, economics, transportation, and social choice [1, 31, 39, 43]. Research has been conducted on problems with single- or multiple-objective optimization under constraints. The main aim of researchers is the development of methods which are able to solve completely non-linear optimization problems. With regard to the difficulties with the constraint function, many methods use some penalty functions to combine the objective functions and constraint function by a penalty function [36, 37, 41, 42]. That allows to transform the initial problem formulation into a single objective function optimization without constraints. For this last formulation, we find many proposed methods to reach the optimal solutions.

We have many studies using the penalty functions in order to make easy the obtaining of optimal solutions but the Lagrangian penalty function is one of the well knew and used in the resolution of constrained optimization problems. Many are works, in the literature, in with Lagrangian penalty function has proved its performance in helping to reach optimal solution in the single-objective optimization. Birgin et al. proposed some results on this topic: global minimization using an augmented Lagrangian method with variable lower-level constraints [5]; practical augmented Lagrangian methods for constrained optimization [4]; optimality properties of an augmented

*Correspondence to: Kounhinir Somé (Email: sokous11@gmail.com). Département de Mathématiques, Laboratoire de Mathématiques, Informatique et Applications (L@MIA), Université Norbert Zongo, BP 376 Koudougou, Burkina Faso.

Lagrangian method on infeasible problems[3]; and a study on the application of an augmented Lagrangian algorithm to some portfolio problems [2]. Cao et al. [7] proposed some work on an augmented Lagrangian interior-point approach to large-scale NLP problems in graphics processing units. Some studies on the convergence of inexact augmented Lagrangian methods for problems with convex constraints are presented by [25]. He et al. [28] are focused on a nonlinear augmented Lagrangian for constrained minimax problems. Augmented Lagrangian applied to the convex quadratics problem is the paper of [32]. Pan et al. [34] used Lagrangian relaxation for the multiple constrained robust shortest path problem. A new augmented Lagrangian method for equality constrained optimization with simple unconstrained sub-problem is the contribution of Zhang et al.

In spite of the abundance of research in the field of single-objective optimization, there is limited work on multiobjective optimization using the Augmented Lagrangian method. In this context, the Augmented Lagrangian method has also attracted the interest of researchers in multiobjective optimization. For instance, Fazzio, N.S. [20] dedicated their thesis to the study of the Augmented Lagrangian method for multiobjective optimization. Another approach was explored in 2020 by G. Cocchi et al. [10], and later refined by the same authors in 2021 in [9]. They propose an enhancement of the previous multiobjective Augmented Lagrangian method by using a set of mutually non-dominant points and employing an Armijo-type line search to enrich the approximate Pareto front. More recently, in 2023, Ashutosh Upadhyay et al. [45] proposed an Augmented Lagrangian method based on the cone method for solving multiobjective optimization problems.

In this work, the main contributions and highlights of this study are presented as follows:

- Introduction of an algorithm developed using the ϵ -constraint approach in conjunction with the augmented Lagrangian method to solve continuous multi-objective problems with convex constraints.
- Analysis of the global convergence of the point sequences generated by the algorithm towards Pareto-stationarity under key assumptions such as that the problem is of a convex nature and the objective functions as well as the constraints are continuous and differentiable.
- Evaluation of the performance of the proposed method compared to state-of-the-art approaches already established in the literature.

To assess the performance of the proposed method, we compare the obtained numerical solutions to those generated by methods such as NSGA-II [12], and BoostDMS [16], using various performance metrics. The obtained results demonstrate that our method outperforms most other methods on the majority of test problems.

The rest of this paper is organized as follows. In Section 2, some preliminaries are given. Section 3 introduces the augmented Lagrange penalty method based on the ϵ -Constraint approach. In Section 4, we present an algorithm for its application accompanied by a theoretical convergence study of the algorithm. In Section 5, we propose the numerical application to test problems existing in the literature. We end in section 6 with a conclusion and some remarks.

2. Preliminaries

As a minimization problem can be transformed into a maximization problem and vice versa, we will present our results for the case of minimization. Multiobjective optimization problems can be formulated by :

$$\begin{aligned} \min : & F(x) = (f_1(x), f_2(x), \dots, f_q(x)) \\ \text{s.t.} : & \begin{cases} g_i(x) \leq 0 & \forall i \in I = 1, \dots, l, \\ x \in \mathbb{R}^n, \end{cases} \end{aligned} \quad (1)$$

where, $F : \mathbb{R}^n \rightarrow \mathbb{R}^q$, $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ are continuous and differentiable functions. $\mathcal{X}$ represents a non-empty convex set. Let $\mathcal{X}$ denote the feasible space of problem (1), defined as $\mathcal{X} = \{x \in \mathbb{R}^n : g(x) \leq 0\}$.

For the rest of this work, we will consider problem (1), whose objective functions are convex and constraints functions are continuous.

The following definitions present the basic concepts of optimal solutions in multiobjective programming.

Definition 1 ([18])

A feasible solution $x^* \in \mathcal{X}$ is called efficient or Pareto optimal, if there is no other $x \in \mathcal{X}$ such that $F(x) \leq F(x^*)$. If x^* is efficient, $F(x^*)$ is called a nondominated point.

Definition 2 ([18])

A feasible solution $x^* \in \mathcal{X}$ is called weakly efficient (weakly Pareto optimal) if there is no $x \in \mathcal{X}$ such that $F(x) < F(x^*)$, i.e., $f_i(x) < f_i(x^*)$ for all $i = 1, 2, \dots, q$.

Now, a necessary condition for a Pareto-stationary point is given by the following definition.

Definition 3

A point $x^* \in \mathcal{X}$ is said to be Pareto-stationary for Problem (1) if, for all $x \in \mathcal{X}$,

$$\max_{j=1,q} \nabla f_j(x^*)^\top (x - x^*) \geq 0.$$

Note also that, if x^* is not Pareto stationary, there exists a $x \in \mathcal{X}$ such as $\max_{j=1,q} \nabla f_j(x^*)^\top (x - x^*) < 0$.

Thus, a well-known equivalent characterization of a Pareto-stationary point from the point of view of the projection is given by the following lemma.

Lemma 1

A point $x^* \in \mathcal{X}$ is said to be Pareto-stationary for Problem (1) if, for all $x \in \mathcal{X}$,

$$x^* = \Pi_{\mathcal{X}} \left[x^* - \max_{j=1,q} \nabla f_j(x^*)^\top (x - x^*) \right], \quad (2)$$

where $\Pi_{\mathcal{X}}[x]$ is the projection operator of the point x in the convex set $\mathcal{X}$.

Now, what should be noted is that if we have one objective, a point x^* is said to be Pareto stationary if $\nabla f(x^*)^\top (x - x^*) \geq 0, \forall x \in \mathcal{X}$, and its equivalent term from the point of view of the projection is given by the following relation $x^* = \Pi_{\mathcal{X}}[x^* - \nabla f(x^*)]$.

Definition 4

We call the ideal point of the problem (1) the vectors $z^* \in \mathbb{R}^q$ whose components z_j^* are obtained by individually minimizing each objective function $f_j, j = 1, 2, \dots, q$ under the constraints of the problem (1). That is to say :

$$\begin{aligned} z_j^* &= \min f_j(x) \\ \text{s.t. : } &\begin{cases} g_i(x) \leq 0, \forall i = 1, \dots, m, \\ x \in \mathbb{R}^n. \end{cases} \end{aligned} \quad (3)$$

The ϵ -constraint approach is a method used to transform problem (1) into a single-objective form as follows:

$$\begin{aligned} \min & f_p(x), \\ \text{s.t. : } &\begin{cases} f_j(x) \leq \epsilon_j, \quad \forall j = 1, 2, \dots, q, \quad j \neq p, \\ g_i(x) \leq 0, \quad \forall i \in I = 1, \dots, l, \\ x \in \mathcal{X}. \end{cases} \end{aligned} \quad (4)$$

where $\epsilon_j \in [\epsilon_j^{\min}, \epsilon_j^{\max}]$ are determined respectively by minimize and maximize $f_j(x)$ under the constraint $x \in \mathcal{X}$. Let us set $g_i(x) = f_i(x) - \epsilon_i, i = 1, \dots, q, i \neq p$, then problem (1) is reworded as follows:

$$\begin{aligned} \min & f_p^\epsilon(x), \\ \text{s.t. : } &\begin{cases} g_i(x) \leq 0, \quad \forall i \in I = 1, \dots, m, \quad i \neq p, \\ x \in \mathcal{X}, \end{cases} \end{aligned} \quad (5)$$

with $m = p - 1 + l$

Proposition 1 ([13])

Any solution to the problem (5) is an optimal Pareto solution for any upper bound vector given by:

$$\epsilon = (\epsilon_1, \epsilon_2, \dots, \epsilon_{p-1}, \epsilon_{p+1}, \epsilon_{p+2}, \dots, \epsilon_q)^\top.$$

3. Main results

The method we have presented in this paper is based on the projected gradient, which requires some conditions before use. It concerns the convexity and compactness of decision space; objective and constraint functions must be differentiable. The above two assumptions provide the basis for these requirements.

Assumption 1

The set $\mathcal{X} \subseteq \mathbb{R}^n$ is closed and convex.

Assumption 2

The objective function F has bounded level sets in the multiobjective sense, i.e., the set $\{x \in \mathbb{R}^n, F(x) \leq F(x_0)\}$ is compact.

3.1. Principle

As a reminder, the Augmented Lagrangian method developed in the work of E. G. Birgin [4] is a method that allows the transformation of a constrained optimization problem into an unconstrained problem. Let us consider problem (5) and pose

$$\mathcal{L}_{\rho_k}(x, \mu^k, \epsilon) = f_p^\epsilon(x) + \frac{\rho_k}{2} \sum_{i=1}^m \max \left\{ 0, g_i(x) + \frac{\mu_i^k}{\rho_k} \right\}^2. \quad (6)$$

The problem (5) after penalization becomes:

$$\min_{x \in \mathcal{X}} \mathcal{L}_{\rho_k}(x, \mu, \epsilon), \quad (7)$$

where $\rho_k > 0$ is a penalization parameter, and $\mu^k \in \mathbb{R}_+^m$ is a vector of the approximate Lagrange multiplier associated with constraints. These parameters are updated at each iteration as follows [25] :

$$\mu_i^{k+1} = \max \{ 0, \mu_i^k + \rho_k g_i(x^{k+1}) \}. \quad (8)$$

Under the assumption that the objective functions and constraints are differentiable, this implies that the function defined by equation (6) is also differentiable. Consequently, the gradient of the function (6) can be expressed as follows:

$$\nabla \mathcal{L}_{\rho_k}(x, \mu^k, \epsilon) = \nabla f_p^\epsilon(x) + \sum_{i=1}^m \left[\max \{ 0, \rho_k g_i(x) + \mu_i^k \} \right] \nabla g_i(x). \quad (9)$$

However, a stationary point of the Augmented Lagrangian sub-problem is given by the following definitions.

Definition 5

Let be ϵ fixed. A point $x^* \in \mathcal{X}$ is a Pareto-stationary point if and only if

$$x^* = \Pi_{\mathcal{X}} [x^* - \nabla \mathcal{L}_{\rho_k}(x, \mu^k, \epsilon)], \quad (10)$$

where $\Pi_{\mathcal{X}}(x)$ is the projection of x in the convex space $\mathcal{X}$.

Thus, the following definition gives a characterization of the α -stationary of the Augmented Lagrangian sub-problem [25].

Definition 6

Let $\alpha \geq 0$ and ϵ . A point $x^* \in \mathcal{X}$ is said to be α -Pareto-stationary of the Lagrange sub-problem if and only if

$$\|x^* - \Pi_{\mathcal{X}} [x^* - \nabla \mathcal{L}_{\rho_k}(x, \mu^k, \epsilon)]\| \leq \alpha. \quad (11)$$

3.2. Algorithm

The following Algorithm presents the augmented Lagrangian method based on the approach of the ϵ -constraint.

Algorithm 1 Augmented Lagrangian based on ϵ -Constraint approach (ALB $_{\epsilon}$) .

- 1: **Require:** $\bar{\mu} \gg 0, \beta > 0, \mu^0 \in \mathbb{R}_+^m, \rho_0 > 0, x^0 \in \mathbb{R}^n, \gamma \geq 0$
- 2: **for** $k = 1, 2, 3, \dots$ **do**
- 3: Find x^{k+1} an α_k -Pareto-Point of $\mathbb{R}^n$ of

$$f_p^\epsilon(x) + \frac{\rho_k}{2} \left[\sum_{i=1}^m \max \left\{ 0, g_i(x) + \frac{\mu_i^k}{\rho_k} \right\}^2 \right]$$

- 4: **for** $i = 1, 2, 3, \dots, m$ **do**
 - 5: $\mu_i^{k+1} = \Pi_{[0; \bar{\mu}]} \left(\max \left\{ 0, \frac{\mu_i^k}{\rho_k} + g_i(x^{k+1}) \right\} \right)$
 - 6: $\sigma_i^{k+1} = \min \left\{ \frac{\mu_i^k}{\rho_k}, -g_i(x^{k+1}) \right\}$
 - 7: **end for**
 - 8: **if** $\|\sigma^{k+1}\| \geq \beta \|\sigma^k\|$ **then**
 - 9: $\rho_{k+1} = \gamma \rho_k$
 - 10: **else**
 - 11: $\rho_{k+1} = \rho_k$
 - 12: **end if**
 - 13: **end for**
-

Algorithm 1 starts with the selection of a priority objective function that will be minimized. The other objective functions are individually minimized under the initial constraints to obtain the components of the vector ϵ . With that, each objective function is transformed into a new constraint by considering the corresponding ϵ value as its maximum value. After that, the initial problem is transformed into a parametric single-objective optimization problem, as shown in equation (5). Next, an arbitrary choice of $\bar{\mu}$ is made to fix the lower and upper bounds of the value of μ . Also, in the same way, μ^0 and ρ_0 are chosen, they will be updated at each iteration in accordance with steps 8 to 11 and at step 13 of the Algorithm 1. Step 7 is dedicated to solving the Lagrangian sub-problem.

Under the assumptions presented above, we apply the projected gradient method [17, 19, 23, 24, 44], which is one of the iterative methods for solving the sub-problem of the Algorithm 1. The determination of step length in the iterative step is determined using Armijo's rule [21, 26]. The well-defined nature of the algorithm follows directly from lemma 4 in the work of Fliege and Sviater [23]. Note that a backtracking Armijo-type line search, is an $\mathbb{R}^n$ -decreasing method, i.e., the values of the objective function always decrease in partial order per component in a finite number of iterations.

3.3. Convergence Study

The following propositions present the admissibility and optimality of the solutions given by the Algorithm 1. Proposition 2 shows that an iterated x^{k+1} of the Algorithm 1 is admissible, i.e., that x^{k+1} belongs to the space $\mathcal{X} = \{x \in \mathbb{R}^n \mid g(x) \leq 0\}$, the Proposition 3 shows that a limit point x^* of the sequence $\{x^k\}$ generated by the Algorithm 1 is weakly Pareto optimal of the problem (5).

Lemma 2

Let x^* be a limit point of the sequence $\{x^{k+1}\}$ generated by the Algorithm 1. If $x^* \in \mathcal{X}$, then for any $i = 1, \dots, m$ such as $g_i(x^*) \leq 0$, we have $\lim_{k \rightarrow \infty} \min \left\{ \frac{\mu_i^k}{\rho_k}, -g_i(x^{k+1}) \right\} = 0$.

Proof

From the Algorithm instructions, we have $\mu_i^k \geq 0$ for any k . There are two possible cases that arise.

(a) ρ_k bounded.

In this case, there is a k_0 such as for any $k \geq k_0$, we have $\rho_k \rightarrow \rho_{k_0}$. Thus, for $k_0 \geq 1$

$$\left\| \min \left\{ \frac{\mu_i^{k_0}}{\rho_{k_0}}, -g(x^{k_0+1}) \right\} \right\| \leq \beta \left\| \min \left\{ \frac{\mu_i^{k_0-1}}{\rho_{k_0-1}}, -g(x^{k_0}) \right\} \right\| \quad (12)$$

$$\left\| \min \left\{ \frac{\mu_i^{k_0+1}}{\rho_{k_0+1}}, -g(x^{k_0+2}) \right\} \right\| \leq \beta \left\| \min \left\{ \frac{\mu_i^{k_0}}{\rho_{k_0}}, -g(x^{k_0+1}) \right\} \right\| \quad (13)$$

$$\left\| \min \left\{ \frac{\mu_i^{k_0+2}}{\rho_{k_0+2}}, -g(x^{k_0+3}) \right\} \right\| \leq \beta^2 \left\| \min \left\{ \frac{\mu_i^{k_0-1}}{\rho_{k_0-1}}, -g(x^{k_0}) \right\} \right\| \quad (14)$$

$$\begin{aligned} & \vdots \\ & \left\| \min \left\{ \frac{\mu_i^{k_0+\eta}}{\rho_{k_0+\eta}}, -g(x^{k_0+\eta+1}) \right\} \right\| \leq \beta^\eta \left\| \min \left\{ \frac{\mu_i^{k_0-1}}{\rho_{k_0-1}}, -g(x^{k_0}) \right\} \right\|. \end{aligned} \quad (15)$$

When $\eta \rightarrow \infty$, since $\beta \in [0; 1[$, we have :

$$\beta^\eta \left\| \min \left\{ \frac{\mu_i^{k_0-1}}{\rho_{k_0-1}}, -g(x^{k_0}) \right\} \right\| \rightarrow 0, \quad (16)$$

which implies that $\left\| \min \left\{ \frac{\mu_i^{k_0}}{\rho_{k_0}}, -g(x^*) \right\} \right\| = 0$. Moreover, if $g_i(x^*) < 0$ we have $\mu_i^* = 0$.

(b) ρ_k Unbound.

By definition, we have the sequence $\{\mu_i^k\}$ bounded which implies

$$\lim_{k \rightarrow \infty} \frac{\mu_i^k}{\rho_k} = 0 \text{ and } g_i(x^*) < 0. \quad (17)$$

Thus, for $k \in K$ sufficiently large, we have : $\lim_{k \rightarrow \infty} \min \left\{ \frac{\mu_i^k}{\rho_k}, -g_i(x^{k+1}) \right\} = 0$. Hence the result. $\square$

Proposition 2

Let $\{x^{k+1}\}$ be a sequence generated by the Algorithm 1 whose limit point is x^* . Then, each point x^* is feasible point for the problem (4).

We will then demonstrate that any limit point generated by Algorithm 1 is Pareto-stationary.

Proposition 3

Let $\{x^{k+1}\}$ be a sequence generated by the Algorithm 1 whose limit point is x^* . Then, any point x^* of the sequence $\{x^{k+1}\}$ is Pareto-stationary of the problem (4).

Proof

Let $K \subseteq \{0, 1, \dots\}$ such that

$$\lim_{k \rightarrow \infty, k \in K} x^{k+1} = x^*. \quad (18)$$

From the previous Proposition, we have x^* that is feasible, i.e., $x \in \mathcal{X}$ such that $g_i(x^*) \leq 0, i = 1, \dots, m$. Let us assume by contradiction that there exists $y \in \mathcal{X}$ such that $g_i(y) \leq 0, i = 1, \dots, m$, et $\nabla f_p^\epsilon(x^*)^\top (y - x^*) < 0$.

From the instruction of the Algorithm 1, posing

$$\tilde{H}_i^k = \max \left\{ 0, g_i(x^{k+1}) + \frac{\mu_i^k}{\rho_k} \right\} \quad (19)$$

and

$$\hat{x}^{k+1} = \Pi_{\mathcal{X}} \left[x^{k+1} - \nabla f_p^\epsilon(x^{k+1}) - \rho_k \sum_{i=1}^m \tilde{H}_i^k \nabla g_i(x^{k+1}) \right] \quad (20)$$

which implies that at each iteration,

$$\|x^{k+1} - \hat{x}^{k+1}\| \leq \alpha_k. \quad (21)$$

From the properties of the projection, we obtain for all $y \in \mathcal{X}$,

$$\left(x^{k+1} - \nabla f_p^\epsilon(x^{k+1}) - \rho_k \sum_{i=1}^m \tilde{H}_i^k \nabla g_i(x^{k+1}) - \hat{x}^{k+1} \right)^\top (y - \hat{x}^{k+1}) \leq 0. \quad (22)$$

By adding and subtracting x^{k+1} , and rearranging, we get

$$\begin{aligned} \nabla f_p^\epsilon(x^{k+1})^\top (y - x^{k+1}) &\geq -\nabla f_p^\epsilon(x^{k+1})^\top (x^{k+1} - \hat{x}^{k+1}) + \|x^{k+1} - \hat{x}^{k+1}\|^2 + \\ &\quad (x^{k+1} - \hat{x}^{k+1})^\top (y - x^{k+1}) - \rho_k \left(\sum_{i=1}^m \tilde{H}_i^k \nabla g_i(x^{k+1}) \right)^\top (y - x^{k+1}) - \\ &\quad \rho_k \left(\sum_{i=1}^m \tilde{H}_i^k \nabla g_i(x^{k+1}) \right)^\top (x^{k+1} - \hat{x}^{k+1}). \end{aligned} \quad (23)$$

Using the convexity of g , the last two terms of (23) can be bounded as follows :

$$\begin{aligned} &-\rho_k \left(\sum_{i=1}^m \tilde{H}_i^k \nabla g_i(x^{k+1}) \right)^\top (y - x^{k+1}) - \rho_k \left(\sum_{i=1}^m \tilde{H}_i^k \nabla g_i(x^{k+1}) \right)^\top (x^{k+1} - \hat{x}^{k+1}) \\ &\geq \rho_k \sum_{i=1}^m \tilde{H}_i^k (g_i(x^{k+1}) - g_i(y)) - \rho_k \left(\sum_{i=1}^m \tilde{H}_i^k \nabla g_i(x^{k+1}) \right)^\top (x^{k+1} - \hat{x}^{k+1}) \geq \\ &\quad \rho_k \sum_{i=1}^m \tilde{H}_i^k g_i(x^{k+1}) - \rho_k \sum_{i=1}^m \tilde{H}_i^k \|\nabla g_i(x^{k+1})\| \|x^{k+1} - \hat{x}^{k+1}\|. \end{aligned} \quad (24)$$

Now, considering the term

$$\rho_k \sum_{i=1}^m \tilde{H}_i^k g_i(x^{k+1}), \text{ if } g_i(x^{k+1}) \geq 0, \quad (25)$$

we have

$$\begin{aligned} \rho_k \sum_{i=1}^m \tilde{H}_i^k g_i(x^{k+1}) &= \sum_{i=1}^m \max \{ 0, \rho_k g_i(x^{k+1}) + \mu_i^k \} g_i(x^{k+1}) \\ &= \sum_{i=1}^m \max \{ 0, \rho_k g_i(x^{k+1})^2 + \mu_i^k g_i(x^{k+1}) \} \\ &\geq \sum_{i=1}^m \max \{ 0, \mu_i^k g_i(x^{k+1}) \} \\ &\geq \sum_{i=1}^m \min \{ 0, \mu_i^k g_i(x^{k+1}) \} \end{aligned} \quad (26)$$

Otherwise, if $g_i(x^{k+1}) \geq 0$ we have

$$\begin{aligned} \rho_k \sum_{i=1}^m \tilde{H}_i^k g_i(x^{k+1}) &= \sum_{i=1}^m \max \{0, \rho_k g_i(x^{k+1}) + \mu_i^k\} g_i(x^{k+1}) \\ &= \sum_{i=1}^m \max \{0, \rho_k g_i(x^{k+1})^2 + \mu_i^k g_i(x^{k+1})\} \\ &\geq \sum_{i=1}^m \min \{0, \mu_i^k g_i(x^{k+1})\} \end{aligned} \quad (27)$$

However, we can rewrite

$$\rho_k \sum_{i=1}^m \tilde{H}_i^k g_i(x^{k+1}) \geq \sum_{i=1}^m \min \{0, \mu_i^k g_i(x^{k+1})\}. \quad (28)$$

Recalling that $\|x^{k+1} - \hat{x}^{k+1}\| \leq \alpha_k$, we can rewrite

$$\begin{aligned} -\rho_k \sum_{i=1}^m \tilde{H}_i^k \|\nabla g_i(x^{k+1})\| \|x^{k+1} - \hat{x}^{k+1}\| &\geq -\rho_k \sum_{i=1}^m \tilde{H}_i^k \|\nabla g_i(x^{k+1})\| \alpha_k \\ &= -\alpha_k \sum_{i=1}^m \max \{0, \rho_k g_i(x^{k+1}) + \mu_i^k\} \|\nabla g_i(x^{k+1})\|. \end{aligned} \quad (29)$$

Equation (23) can therefore be rewritten as follows

$$\begin{aligned} \nabla f_p^\epsilon(x^{k+1})^\top (y - x^{k+1}) &\geq -\nabla f_p^\epsilon(x^{k+1})^\top (x^{k+1} - \hat{x}^{k+1}) + \|x^{k+1} - \hat{x}^{k+1}\|^2 + \\ &\quad (x^{k+1} - \hat{x}^{k+1})^\top (y - x^{k+1}) \sum_{i=1}^m \min \{0, \mu_i^k g_i(x^{k+1})\} \\ &\quad - \alpha_k \sum_{i=1}^m \max \{0, \rho_k g_i(x^{k+1}) + \mu_i^k\} \|\nabla g_i(x^{k+1})\| \end{aligned} \quad (30)$$

Now, there are two possible cases :

- (i) ρ_k bounded,
according to Lemma 2, we have $\mu_i^k = 0$
- (ii) $\rho_k \rightarrow \infty$ there exists a k_0 such that for all $k \geq k_0$, we obtain

$$\rho_k g_i(x^{k+1}) + \mu_i^k < 0 \text{ and } \max \{0, \rho_k g_i(x^{k+1}) + \mu_i^k\} = 0. \quad (31)$$

In each case (i) and (ii), since $g(x^*) \leq 0$, for k sufficiently large, $k \in K$ and continuous of ∇f_p^ϵ and ∇g , with $\|x^{k+1} - \hat{x}^{k+1}\| \rightarrow 0$, $\alpha_k \rightarrow 0$ and $\sum_{i=1}^m \min \{0, \mu_i^k g_i(x^{k+1})\} = 0$ we obtain

$$\nabla f_p^\epsilon(x^*)^\top (y - x^*) \geq 0. \quad (32)$$

This contradicts the initial hypothesis. □

4. Numerical Experiences

In this section, we present some numerical results of ALB_ϵ on some test problems. In addition, a comparison is done with two other methods namely NSGA-II and BoostDMS method.

The following Table 1 gives the used parameters of each method.

Table 1. Value of used parameters for methods

Methods Parameters	ALB_ϵ	NSGA-II	BoostDMS
Population	100	100	Parameter par default
Iteration number	100	2000	Parameter par default
Number of objective function assessment	10 000	200 000	10 000

In total, we performed the simulation on 50 test problems recorded in Table 2. Among the test problems found in the literature, those that do not have boundary constraints, we have defined research boundaries on these problems. For the implementation of ALB_ϵ , we used the parameters $\lambda_0 = (0, 1, 1, \dots)^\top \in \mathbb{R}^m$, $\rho_0 = 1$, $\alpha^* = 10^{-6}$, $\sigma = 0.1$, $\gamma = 4$.

Table 2. List of Multiobjective optimization test problems

Problems	n	q	Parameters bornes	Source
BNH1	2	2	$[0, 5]^2$	[29]
SCH	1	2	$[-4, 4]$	[37]
ZDT1	a	2	$[0, 1]^n$	[12]
LAP1	2	2	$[-7, 1]^2$	[10]
LAP2	b	2	$[0, 1]^n$	[10]
JOS1	c	2	$[0, 1]^n$	[29]
ZLT1	d	2	$[-1, 1]^n$	[29]
MLF1	1	2	$[0, 20]$	[29]
BK1	2	2	$[-5, 10]^2$	[29]
SSFYY1	1	2	$[-100, 100]$	[29]
VU2	2	2	$[-3, 3]^2$	[29]
COSH	1	2	$[0, 5]$	[48]
KD1	2	2	$[0.1, 1] \times [0, 1]$	[14]
SRN	2	2	$[-20, 20]^2$	[29]
BNH3	2	2	$[-7, 4]^2$	[29]
SSFYY2	2	2	$[-100, 100]^2$	[29]
MIN-Ex	2	2	$[0.1, 1] \times [0, 5]$	[37]

We have dealt with the test problems with :

$a \in \{2, 5, 10, 15, 20, 25, 30\}$; $b \in \{2, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100\}$; $c \in \{2, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50\}$; $d \in \{30, 35, 40, 45, 50\}$.

As in [6, 11, 38], we used the same metrics proposed for the comparison of the Algorithms. The metrics used are the purity metric, the spreads metric, and the hyper-volume indicator. The purity metric measures the quality of the Pareto front generated by an Algorithm. It gives the percentage of non-dominated solutions generated by method [11]. The purity metric is given by the following formula :

$$Purity(S) = \frac{|F_{p,s} \cap F_p|}{|F_{p,s}|}, \quad (33)$$

with $F_{p,s}$ the solutions generated by a solver $s \in S$ for a problem $p \in P$ where S is the set of solvers and P the set of test problems. F_p represents the set of solutions generated by all solvers for the problem p ($F_p = \bigcup F_{p,s}$) without the dominated points.

The spread metrics used are Γ -Spread and Δ -Spread. The Γ -Spread metric measures the maximum spacing of solutions generated by a solver [11]. It is given by the following formula :

$$\Gamma\text{-Spread}(S) = \max_{j \in \{1, \dots, q\}} \left(\max_{i \in \{0, \dots, N\}} \{\delta_{i,j}\} \right), \quad (34)$$

where N represents the number of solutions generated by a solver, m is the number of objective functions, and $\delta_{i,j} = f_j(x_{k+1}) - f_j(x_k)$ whose values of $f_j(x_k)$ are arranged in ascending order. The Δ -Spread metric measures the distribution of solutions generated by a solver [11]. It is calculated by the following formula :

$$\Delta\text{-Spread}(S) = \max_{j \in \{1, \dots, q\}} \frac{\delta_{0,j} + \delta_{N,j} + \sum_{i=1}^{N-1} |\delta_{i,j} - \bar{\delta}_{i,j}|}{\delta_{0,j} + \delta_{N,j} + (N-1)\bar{\delta}_{i,j}}, \quad (35)$$

where $\bar{\delta}_{i,j}$ is the average of the $\delta_{i,j}$ with $j = 1, \dots, N-1$. $\delta_{0,j}$ and $\delta_{N,j}$ represent the extreme points indexed by 0 and $N+1$. Thus, we used the technique proposed in [11] to compute the extreme points for problems that do not have an analytic front. We first removed the dominant points from the meeting on all these fronts. Then, for each component of the objective function, we selected the pair corresponding to the highest distance in pairs measured using $f_j(\cdot)$.

For a minimization problem, the hyper-volume indicator measures the volume of the part of the objective space that is dominated by the computed approximation to the Pareto front of a problem, bounded by a reference point $W^p \in \mathbb{R}^q$ [22]. The following formula gives the hyper-volume indicator :

where $Volume(\cdot)$ denotes the Lebesgue measure. As in [6], the coordinates of the reference point W^p are determined by the relation

$$w_j^p = \max \left\{ f_j(x) : z \in \bigcup_{s \in S} F_{p,s} \right\}. \quad (36)$$

The scaling of $H(F_{p,s})$ is given by :

$$HV_{p,s} = \frac{H(F_{p,s})}{\prod_{j=1}^m (w_j^p - l_j^p)} \in [0, 1], \quad (37)$$

with $l_j^p = \min \{f_j(x) : z \in \bigcup_{s \in S} F_{p,s}\}$.

We then use the performance profiles proposed in [16, 38, 6] for an appreciation of the performance of the four metrics presented above. We refer the reader to the articles cited above for more information on performance profiles. Recall that the performance profiles are presented by a diagram of a cumulative distribution function $\rho(\tau)$ which is defined as follows :

$$\rho_s(\tau) = \frac{1}{|P|} |\{p \in P : r_{p,s} \leq \tau\}| \quad (38)$$

with $r_{p,s} = \frac{t_{p,s}}{\min \{t_{p,\bar{s}} : \bar{s} \in S\}}$. Since performance profiles are used for metrics whose lowest value indicates better performance, metrics such as Purity and Hypervolume, we will pose $t_{p,s} = 1/t_{p,s}$ as proposed in [6].

4.1. Performance Profiles

Now, we present the performance profiles on the four metrics presented above on the 50 test problems. Figures 1, 2 and 3 represents the performance profile on the purity of ALB ϵ , NSGA-II, and BoostDMS methods.

We can observe that on purity, ALB_{ϵ} is better than NSGA-II and BoostDMS in terms of efficiency. Figures 4,5 and 6 represents the performance profile of the Γ -Spread metric of the three methods. We can observe a dominance of the ALB_{ϵ} method by the NSGA-II and BoostDMS methods. Considering Figures 7,8 and 9 and Figures 10,11 and 12, which represents the performances profiles of the Δ -Spread metric and the Hyper-volume indicator, we notice that the ALB_{ϵ} method remains competitive.





Figure 1. Performance profile of Purity of ALB_{ϵ} and NSGA-II Figure 2. Performance profile of Purity of ALB_{ϵ} and BoostDMS





Figure 3. Performance profile of Purity of ALB_{ϵ} , BoostDMS and NSGA-II Figure 4. Performance profile of Γ -Spread of ALB_{ϵ} and NSGA-II





Figure 5. Performance profile of Γ -Spread of ALB_ϵ and Figure 6. Performance profile of Γ -Spread of ALB_ϵ BoostDMS and NSGA-II





Figure 7. Performance profile of δ -Spread of ALB_ϵ and Figure 8. Performance profile of δ -Spread of ALB_ϵ and BoostDMS





Figure 9. Performance profile of δ -Spread of ALB_ϵ Figure 10. Performance profile of Hypervolume of ALB_ϵ and BoostDMS and NSGA-II

The following tables present a statistical study of the 50 test problems. It is about the average, minimum and maximum values obtained on the four metrics.





Figure 11. Performance profile of Hypervolume of ALB_ϵ and Figure 12. Performance profile of Hypervolume of ALB_ϵ , BoostDMS and NSGA-II

On these four metrics, we notice that the ALB_ϵ method obtains better values on the purity and the Γ -Spread metrics. For the Δ -Spread metric, ALB_ϵ wins over BoostDMS.

Table 3. Statistical study of performance values on Purity

Methods	average	Minimum	Maximum
ALB_ϵ	0.9851	0.5600	1.0000
$NSGA-II$	0.5286	0.0000	1.0000
$BoostDMS$	0.7244	0.0000	1.0000

Table 4. Statistical study of performance values on $\Gamma - Spread$

Methods	average	Minimum	Maximum
ALB_ϵ	8.2902	0.0104	50.4685
$NSGA-II$	16.7456	0.0323	100.0021
$BoostDMS$	11.2006	0.0036	83.6838

Table 5. Statistical study of performance values on Δ -Spread

Methods	average	Minimum	Maximum
ALB_ϵ	1.0153	0.5683	7.1249
$NSGA-II$	0.9477	0.7312	4.4815
$BoostDMS$	1.0834	0.7435	4.9618

4.2. Study of problems with an analytical front

It about the problems SCH , $COSH$, $MIN - Ex$, $ZDT1$ (with 30 variables) and KD_1 . We use these problems for a comparison of the four metrics against the true front. For the KD_1 problem the global minimum is obtained

Table 6. Statistical study of performance values on Hypervolume

Methods	average	Minimum	Maximum
ALB_ϵ	0.7012	0.3832	0.8987
$NSGA-II$	0.7066	0.3858	0.8985
$BoostDMS$	0.7166	0.3883	0.8985

for $x_2 \approx 0.2$ and the local minimum for $x_2 \approx 0.6$ with $g(0.2) = 0.7057$ and $g(0.6) = 1.2$ [14]. In this work, we use the case $x_2 \approx 0.2$. The same number of solutions for the three methods are imposed. The values obtained for these test problems are recorded in the Tables 7, 8, 9, 10. The Figures 13,14,15,16,17 represent the Pareto fronts obtained. We see that except for the bimodal problem $KD1$, the ALB_ϵ method wins over the two methods on the purity metric. For the Γ -Spread metric, the ALB_ϵ method is better on the Min-Ex, Cosh, $KD1$, and $ZDT1$. For the Δ -Spread metric, only the $ZDT1$ problem is won by the ALB_ϵ method. For hypervolume, ALB_ϵ is competitive on both Algorithm.




Figure 13. Pareto front SCH




Figure 14. Pareto front of COSH




Figure 15. Pareto front Min-Ex




Figure 16. Pareto front of ZDT1




Figure 17. Pareto front of KD1

Table 7. Value on performance measures of Purity

Problems	Min-Ex	COSH	KD ₁	SCH	ZDT ₁
ALB_{ϵ}	1.0000	1.0000	0.8424	0.9993	1.0000
$NSGA-II$	1.0000	1.0000	0.9799	0.9993	0.9159
$BoostDMS$	0.8788	1.0000	0.7908	0.9993	1.0000

Table 8. Value on performance measures of $\Gamma - Spread$

Problems	Min-Ex	Cosh	KD ₁	SCH	ZDT ₁
ALB_{ϵ}	0.0104	0.7901	0.0378	0.2134	0.0029
$NSGA-II$	0.0667	0.7901	4.2170	0.0234	1.0797
$BoostDMS$	0.1618	0.7901	0.2023	0.0091	0.0442

Table 9. Value on performance measures of Δ -Spread

Problems	Min-Ex	Cosh	KD ₁	SCH	ZDT ₁
ALB_{ϵ}	1.0175	1.1477	1.0647	1.0000	0.7500
$NSGA-II$	0.8451	0.9880	0.9951	0.8343	1.0978
$BoostDMS$	0.9714	1.1145	1.0423	0.7638	0.7579

Table 10. Value on performance measures of Hyper-volume

Problems	Min-Ex	Cosh	KD ₁	SCH	ZDT ₁
ALB_{ϵ}	0.8263	0.8985	0.8246	0.8330	0.6659
$NSGA-II$	0.8263	0.8985	0.8987	0.8330	0.8570
$BoostDMS$	0.8262	0.8984	0.8304	0.8331	0.6658

4.3. Comment

From the theoretical and numerical results of the ALB_ϵ method, we can say that it is competitive in solving multiobjective optimization problems since it gives better results on the generation of non-dominated solutions. For the maximum spacing, we can say that it is due to the ϵ -constraint approach. Since we choose a priority function, this can affect the maximum spacing of solutions. For the diversity of solutions and the hypervolume metric, we observe that the ALB_ϵ method remains competitive.

5. Conclusion

In this work, an augmented Lagrangian function was combined with an ϵ -constraint approach to solve constrained multiobjective optimization problems. It consists of transforming constrained multiobjective optimization problems into unconstrained single objective optimization problems. The proposed algorithm is deterministic, and the Pareto optimality of the provided solutions has been justified. First, we present the algorithm of the method and some theoretical convergence results through some propositions. Then, we showed the numerical results of fifty test problems from the literature that we solved with our method, NSGA-II and BoostDMS, which are well-known and used methods. Finally, we compared the numerical performance of these three methods. This study has been focused on convergence and distribution parameters using obtained solutions. According to this study, it appears that ALB_ϵ is the best choice for the resolution of multiobjective optimization problems when the admissible solutions set is bounded and convex, and the objective space is compact.

The next phase of our research will focus on the extension of the algorithm to tackle non-convex objective functions, on the one hand, and on the other, tackle non-differentiable cases. Furthermore, we are interested in the use of this method to solve real-life problems.

Authors contributions

All authors contributed equally to this work.

Funding

This study is not funded.

Data availability

No underlying data was collected or produced in this study.

Declarations

Ethics approval

Not applicable.

Conflict of interest

The authors declare no competing interest.

REFERENCES

1. Bhatia, M. & Ran, A. Integrated Farm Model for Optimal Allocation of Resources- A Linear Programming Approach. *Applications And Applied Mathematics: An International Journal (AAM)*. **15**, Article 37 (2020)
2. Birgin, E. & Martínez, J. On the Application of an Augmented Lagrangian Algorithm to Some Portfolio Problems. *EURO Journal On Computational Optimization*. **4**, 79-92 (2016)
3. Birgin, E., Martínez, J. & Prudente, L. Optimality Properties of an Augmented Lagrangian Method on Infeasible Problems. *Computational Optimization And Applications*. **60**, 609-631 (2015)
4. Birgin, E. & Martínez, J. Practical Augmented Lagrangian Methods for Constrained Optimization. *Encyclopedia Of Optimization*. pp. 3013-3023 (2014)
5. Birgin, E., Floudas, C. & Martínez, J. Global Minimization Using an Augmented Lagrangian Method with Variable Lower-Level Constraints. *Mathematical Programming*. **125**, 139-162 (2010)
6. Brás, C. & Custódio, A. On the Use of Polynomial Models in Multiobjective Directional Direct Search. *Computational Optimization And Applications*. **77**, 897-918 (2020)
7. Cao, Y., Seth, A. & Laird, C. An Augmented Lagrangian Interior-Point Approach for Large-Scale NLP Problems on Graphics Processing Units. *Computers & Chemical Engineering*. **85**, 76-83 (2016)
8. Chen, L. & Goldfarb, D. Interior-Point L2-penalty Methods for Nonlinear Programming with Strong Global Convergence Properties. *Mathematical Programming*. **108**, 1-36 (2006)
9. Cocchi, G., Lapucci, M. & Mansueto, P. Pareto Front Approximation through a Multi-Objective Augmented Lagrangian Method. *EURO Journal On Computational Optimization*. **9**, 1-12 (2021)
10. Cocchi, G. & Lapucci, M. An Augmented Lagrangian Algorithm for Multi-Objective Optimization. *Computational Optimization And Applications*. **77**, 29-56 (2020)
11. Custódio, A., Madeira, J., Vaz, A. & Vicente, L. Direct Multisearch for Multiobjective Optimization. *SIAM Journal On Optimization*. **21**, 1109-1140 (2011)
12. Deb, K., Pratap, A., Agarwal, S. & Meyarivan, T. A Fast and Elitist Multiobjective Genetic Algorithm: NSGA-II. *IEEE Transactions On Evolutionary Computation*. **6**, 182-197 (2002)
13. Deb, K. Multi-Objective Optimization Using Evolutionary Algorithms. (John Wiley & Sons Inc, 2001)
14. Deb, K. Multi-Objective Genetic Algorithms: Problem Difficulties and Construction of Test Problems. *Evolutionary Computation*. **7**, 205-230 (1999)
15. De Carvalho, E., Dos Santos, A. & Ma, T. Reduced Gradient Method Combined with Augmented Lagrangian and Barrier for the Optimal Power Flow Problem. *Applied Mathematics And Computation*. **200**, 529-536 (2008)
16. Dolan, E. & Moré, J. Benchmarking Optimization Software with Performance Profiles. *Mathematical Programming*. **91**, 201-213 (2002)
17. Dussault, J. & Fournier, G. On the Convergence of the Projected Gradient Method. *Journal Of Optimization Theory And Applications*. **77**, 197-208 (1993)
18. Ehrgott, M. Multicriteria Optimization. Springer Science & Business Media, (2005).
19. Fan, S. & Yao, Y. Strong Convergence of a Projected Gradient Method. *Journal Of Applied Mathematics*. **2012**, 1-10 (2012)
20. Fazzio, N. Teoría y métodos para problemas de optimización multiobjetivo. (Universidad Nacional de La Plata, 2018).
21. Fitriah, Z. & Anam, S. Modified Armijo Rule on Gradient Descent and Conjugate Gradient. *E-Jurnal Matematika*. **6**, 196-204 (2017)
22. Fleischer, M. The Measure of Pareto Optima Applications to Multi-objective Metaheuristics. *Evolutionary Multi-Criterion Optimization*. pp. 519-533 (2003)
23. Fliege, J. & Svaiter, B. Steepest Descent Methods for Multicriteria Optimization. *Mathematical Methods Of Operations Research (ZOR)*. **51**, 479-494 (2000)
24. Fukuda, E. & Drummond, L. On the Convergence of the Projected Gradient Method for Vector Optimization. *Optimization. A Journal Of Mathematical Programming And Operations Research*. **60**, 1009-1021 (2011)
25. Galvan, G. & Lapucci, M. On the Convergence of Inexact Augmented Lagrangian Methods for Problems with Convex Constraints. *Operations Research Letters*. **47**, 185-189 (2019)
26. García-Palomares, U. & Restuccia, A. Application of the Armijo Stepsize Rule to the Solution of a Nonlinear System of Equalities and Inequalities. *Journal Of Optimization Theory And Applications*. **41**, 405-415 (1983)
27. Hassan, M. & Baharum, A. Generalized Logarithmic Penalty Function Method for Solving Smooth Nonlinear Programming Involving Inex Functions. *Arab Journal Of Basic And Applied Sciences*. **26**, 202-214 (2019)
28. He, S. & Zhou, S. A Nonlinear Augmented Lagrangian for Constrained Minimax Problems. *Applied Mathematics And Computation*. **2018**, 4567-4579 (2011)
29. Huband, S., Hingston, P., Barone, L. & While, L. A Review of Multiobjective Test Problems and a Scalable Test Problem Toolkit. *IEEE Transactions On Evolutionary Computation*. **10**, 477-506 (2006)
30. Kanzow, C. & Steck, D. An example comparing the standard and safeguarded augmented Lagrangian methods. *Operations Research Letters*. **45**, 598-603 (2017)
31. Lian, S., Meng, S. & Wang, Y. An Objective Penalty Function-Based Method for Inequality Constrained Minimization Problem. *Mathematical Problems In Engineering*. **2018**, 1-7 (2018)
32. Marcilio, D. & Matioli, L. Augmented Lagrangian Applied to Convex Quadratic Problems. *Applied Mathematics And Computation*. **200**, 480-485 (2008)
33. Meng, Z., Dang, C. & Yang, X. On the Smoothing of the Square-Root Exact Penalty Function for Inequality Constrained Optimization. *Computational Optimization And Applications*. **35**, 375-398 (2006)
34. Pan, Y. Lagrangian Relaxation for the Multiple Constrained Robust Shortest Path Problem. *Mathematical Problems In Engineering*. **2019**, 1-13 (2019)
35. Rocha, A., Martins, T. & Fernandes, E. An Augmented Lagrangian Fish Swarm Based Method for Global Optimization. *Journal Of Computational And Applied Mathematics*. **235**, 4611-4620 (2011)

36. Som, A., Compaoré, A., Somé, K. & Somé, B. Effect of Aggregation Function in MOMA-Plus Method For Obtaining Pareto Optimal Solutions. *Applications And Applied Mathematics: An International Journal (AAM)*. **17**, Article 14 (2022)
37. Somé, K., Compaoré, A., Sawadogo, W. & Somé, B. Improving of Numerical Performance of The MOMA Method. *International Journal Of Mathematics And Computer Research*. **5**, 1817-1827 (2017)
38. Tavares, S., Brás, C., Custódio, A., Duarte, V. & Medeiros, P. Parallel Strategies for Direct Multisearch. *Numerical Algorithms*. **2022**, 1757-1788 (2022)
39. Temelcan, G., Albayrak, I. & Sivri, M. Solving Multi-Objective Linear Fractional Programming Problems via Zero-Sum Game. *Applications & Applied Mathematics*. **17**, Article 12 (2022)
40. Tian, B., Yang, X. & Meng, K. An Interior-Point L1/2-Penalty Method for Inequality Constrained Nonlinear Optimization. *Journal Of Industrial And Management Optimization*. **12**, 949-973 (2016)
41. Tougma, A., Kaboré, A. & Somé, K. Hyperbolic Augmented Lagrangian algorithm for multiobjective optimization problems. *Gulf Journal Of Mathematics*. **16**, 151-170 (2024)
42. Tougma, A., Some, K. (2024) Inexact Exponential Penalty Function with the Augmented Lagrangian for Multiobjective Optimization Algorithms, *Journal of Applied Mathematics*, vol. 2024, Article ID 9615743, 19 pages.
43. Tougma, A., Yougbaré, W. & Zongo, M. (2021). Performance Analysis of Higher Education Institutions using AHP and PROMETHEE. *AADM*. **27(2)**, 265-284.
44. Tougma, A., Somé, K. & Compaoré, A. (2023). Extension of the Projected Gradient and Armijo's Rule Concepts for Solving Convex Nonlinear Multiobjective Optimization Problems. *Applied Analysis And Optimization*. **7(3)**, 263-278.
45. Upadhayay, A., Ghosh, D., Ansari, Q. & Jauny Augmented Lagrangian Cone Method for Multiobjective Optimization Problems with an Application to an Optimal Control Problem. *Optimization And Engineering*. **24**, 1633-1665 (2022).
46. Wu*, Z., Bai, F., Yang, X. & Zhang, L. An Exact Lower Order Penalty Function and Its Smoothing in Nonlinear Programming. *Optimization. A Journal Of Mathematical Programming And Operations Research*. **53**, 51-68 (2004)
47. Zangwill, W. Non-Linear Programming Via Penalty Functions. *Management Science*. **13**, 344-358 (1967)
48. Zhang, W. & Gao, T. A Min-Max Method with Adaptive Weightings for Uniformly Spaced Pareto Optimum Points. *Computers & Structures*. **84**, 1760-1769 (2006)
49. Zhang, H. & Ni, Q. A New Augmented Lagrangian Method for Equality Constrained Optimization with Simple Unconstrained Subproblem. *Discrete Dynamics In Nature And Society*. **2017**, 1-9 (2017)