



Fixed Point Results in Neutrosophic n-Controlled Bipolar Metric Spaces

M. Rathivel¹, C. Inbam² and M. Jeyaraman^{3,*}

¹Research Scholar, P.G. and Research Department of Mathematics, Raja Doraisingam Govt. Arts College, Sivagangai, Affiliated to Alagappa University, Karaikudi, Tamilnadu, India. Email: rathiravi52379@gmail.com

²Associate Professor, Department of Mathematics, Sri Meenakshi Govt. Arts College For Women(A), Affiliated to Madurai Kamaraj University, Madurai, Tamilnadu, India. Email: cinbam1978@gmail.com

³Associate Professor and Head, P. G. and Research Department of Mathematics, Raja Doraisingam Govt. Arts College, Sivagangai, Affiliated to Alagappa University, Karaikudi, Tamilnadu, India. E-mail: jeya.math@gmail.com.

Abstract We present the idea of Neutrosophic n-Controlled Bipolar Metric space in this study. We use n-non-comparable functions to establish several fixed point results. Furthermore, we employ a version of the Banach contraction principle to extrapolate the conclusions and provide a few non-trivial examples. Subsequently, we apply the main findings to solve financial modeling problems in fractional differential equations.

Keywords Fixed point, Neutrosophic n-controlled bipolar Metric Space, Differential equation.

AMS 2010 subject classifications 47H10, 54H25.

DOI: 10.19139/soic-2310-5070-2945

1. Introduction

Fretchet introduced the concept of metric space in his dissertation from 1906. Later, in 1922, Banach proved the Banach contraction principle in his doctoral dissertation. Since then, several researchers have been experimenting with this idea that have been under numerous circumstances. It is thought to be the most important tool for non-linear analysis. It explains why there is only one fixed point in all contractive mappings in complete metric spaces. It is an extension and generalisation of many metric space types. Zadeh [22] originated the idea of a fuzzy set in 1965. It is a modified form of a conventional set wherein each element has a membership value that goes within a reasonable range. In 1998, Smarandache coined the phrase “neutrosophic set” and proved it next to Sowndrarajan [21]. As a team, they revealed a number of noteworthy findings from neutrosophic metric space. Simsek and Kirisci [13] moved out in 2019 and proposed neutrosophic metric space (*NMS*). Remarkable fixed point results in neutrosophic metric space were validated in 2020 by Sowdrarajan and Jeyaraman et al. [21].

Very recently, in 2016, Mutlu and Gurdal [1] created their concept of bipolar metric spaces (*BMS*). They also evaluated several fixed point conclusions on this space. In the present paper, we will keep trying to study fixed points in the Neutrosophic n-Controlled Bipolar Metric space (*NnCBMS*). Especially some common fixed-point results for an array of contra variants and covariant. Plenty of structures were obtained and the results were standardised across numerous spaces using this topic. It could be achieved to preserve some of the fundamental findings on this topic, such as controlled metric type spaces (*CMS*) and the associated contraction principle in [20], controlled neutrosophic metric spaces (*CNMS*) and certain associated fixed point results in [4], and, more

*Correspondence to: M. Jeyaraman (Email: jeya.math@gmail.com). Associate Professor and Head, P. G. and Research Department of Mathematics, Raja Doraisingam Govt. Arts College, Sivagangai, Affiliated to Alagappa University, karaikudi, Tamilnadu, India.

recently, the unique characteristics of metric spaces in [20].

We establish a number of fixed point findings using n -non-comparable functions. Likewise, we provides several non-trivial cases and extend the results using a variant of the Banach contraction principle. We illustrate the validity and value of the hypothesis based on the results achieved. The ensuing findings corroborate and expand upon the concepts outlined in the paper [20] in a number of afterward literatures.

2. Preliminaries

Definition 2.1. [21] An ordered 6-tuple $(Q, Z, K, Y, \otimes, \odot)$ is called *NMS* if $Q \neq \emptyset$, \otimes *NCTN*, \odot *NCTC* and Z, K, Y are *NS* on $Q \times Q \times (0, \infty)$ the following requirements are met : For all $f, g, h \in Q, \Theta, x > 0$.

- (a) $0 \leq Z(f, g, \Theta) \leq 1; 0 \leq K(f, g, \Theta) \leq 1; 0 \leq Y(f, g, \Theta) \leq 1$;
- (b) $Z(f, g, \Theta) + K(f, g, \Theta) + Y(f, g, \Theta) \leq 3$;
- (c) $Z(f, g, \Theta) = 1, \forall \Theta > 0, \Leftrightarrow f = g$;
- (d) $Z(f, g, \Theta) = Z(g, f, \Theta)$, for $\Theta > 0$;
- (e) $Z(f, g, \Theta) \otimes Z(g, h, x) \leq Z(f, h, \Theta + x) \forall \Theta, x > 0$;
- (f) $Z(f, g, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is *NCTS* and $\lim_{z \rightarrow +\infty} Z(f, g, \Theta) = 1$;
- (g) $K(f, g, \Theta) = 0, \forall \Theta > 0, \Leftrightarrow f = g$;
- (h) $K(f, g, \Theta) = K(f, g, \Theta)$, for $\Theta > 0$;
- (i) $K(f, g, \Theta) \odot K(g, h, x) \geq K(f, h, \Theta + x) \forall \Theta, x > 0$;
- (j) $K(f, g, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is *NCTS* and $\lim_{z \rightarrow +\infty} K(f, g, \Theta) = 0$;
- (k) $Y(f, g, \Theta) = 0, \forall \Theta > 0, \Leftrightarrow f = g$;
- (l) $Y(f, g, \Theta) = Y(\Theta, \epsilon, z)$; for $\Theta > 0$;
- (m) $Y(f, g, \Theta) \odot D(g, h, x) \geq Y(f, h, \Theta + x) \forall \Theta, x > 0$;
- (n) $Y(f, g, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is *NCTS* and $\lim_{z \rightarrow +\infty} Y(f, g, \Theta) = 0$;

Then, $(Q, Z, K, Y, \otimes, \odot)$ is called a *NMS*.

Definition 2.2. [20] Given Υ , let $Q \neq \emptyset$ and $\Upsilon : Q \times Q \rightarrow [1, +\infty)$ are incompetent mapping, if $Z : Q \times Q \rightarrow (0, +\infty)$ is said to be a *CMS* if

- (a) $Z(\epsilon, \Theta) = 0$ iff $\epsilon = \Theta$;
- (b) $Z(\epsilon, \Theta) = Z(\Theta, \epsilon)$;
- (c) $Z(\epsilon, \Theta) \leq \Upsilon(\epsilon, \eta)Z(\epsilon, \eta) + \Upsilon(\eta, \Theta)Z(\eta, \Theta)$; for every $\epsilon, \Theta, \eta \in Q$.

Definition 2.3. [4] Let $Q \neq \emptyset$ and $\Upsilon : Q \times Q \rightarrow [1, +\infty)$, \otimes *NCTN*, \odot *NCTC* and Z, K, Y are *NS* on $Q \times Q \times (0, \infty)$ the following requirements are met: For all $f, g, h \in Q, \Theta, x > 0$

- (a) $0 \leq Z(f, g, \Theta) \leq 1; 0 \leq K(f, g, \Theta) \leq 1; 0 \leq Y(f, g, \Theta) \leq 1$;
- (b) $Z(f, g, \Theta) + K(f, g, \Theta) + Y(f, g, \Theta) \leq 3$;
- (c) $Z(f, g, 0) = 0$;
- (d) $Z(f, g, \Theta) = 1, \forall \Theta > 0, \Leftrightarrow f = g$;
- (e) $Z(f, g, \Theta) = Z(g, f, \Theta)$;
- (f) $Z(f, h, \Theta + x) \geq Z\left(f, g, \frac{\Theta}{\Upsilon(f, g)}\right) \otimes Z\left(g, h, \frac{x}{\Upsilon(g, h)}\right)$;
- (g) $Z(f, g, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is *CTS* and $\lim_{z \rightarrow +\infty} Z(f, g, \Theta) = 1$;
- (h) $K(f, g, 0) = 1$;
- (i) $K(f, g, \Theta) = 0, \forall \Theta > 0 \Leftrightarrow f = g$;
- (j) $K(f, g, \Theta) = K(g, f, \Theta)$;
- (k) $K(f, h, \Theta + x) \leq K\left(f, g, \frac{\Theta}{\Upsilon(f, g)}\right) \odot K\left(g, h, \frac{x}{\Upsilon(g, h)}\right)$;
- (l) $K(f, g, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is *CTS* and $\lim_{z \rightarrow +\infty} K(f, g, \Theta) = 0$;
- (m) $Y(f, g, 0) = 1$;

- (n) $Y(f, g, \Theta) = 0, \forall \Theta > 0 \Leftrightarrow f = g$;
 (o) $Y(f, g, \Theta) = Y(g, f, \Theta)$;
 (p) $Y(f, h, \Theta + x) \leq Y\left(f, g, \frac{\Theta}{\Upsilon(f, g)}\right) \odot Y\left(g, h, \frac{x}{\Upsilon(g, h)}\right)$;
 (q) $Y(f, g, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is *CTS* and $\lim_{z \rightarrow +\infty} Y(f, g, \Theta) = 0$;

Then, $(Q, Z, K, Y, \otimes, \odot)$ is called a *NCMS*.

Definition 2.4. [9] Suppose $Q, L \neq \emptyset$ and $\Upsilon : Q \times Q \rightarrow [1, +\infty)$ are considered as a incompetant mappings, \otimes as *CTN* defined as $r \otimes s = \min\{r, s\}$ and \odot as *CTC* defined as $r \odot s = \max\{r, s\}$ and Z, K, Y are *NS* on $Q \times L \times (0, +\infty)$ is characterized *NBMS*, if for each one $(Q, L, Z, K, Y, \otimes, \odot)$ fulfills all $f \in Q, l \in L$ and $\Theta, x, z > 0$ holds the following:

- (a) $0 \leq Z(f, g, \Theta) \leq 1; 0 \leq K(f, g, \Theta) \leq 1; 0 \leq Y(f, g, \Theta) \leq 1$;
 (b) $Z(f, g, \Theta) + K(f, g, \Theta) + Y(f, g, \Theta) \leq 3$;
 (c) $Z(f, g, 0) > 0$, for every value $f, g \in Q \times L$;
 (d) $Z(f, g, \Theta) = 1$ iff $f = l$ for $f \in Q, g \in L$;
 (e) $Z(f, g, \Theta) = Z(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
 (f) $Z(f_1, g_2, \Theta + x + z) \geq Z(f_1, g_1, \Theta) \otimes Z(f_2, g_1, x) \otimes Z(f_2, g_2, z)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L \forall \Theta, x, z > 0$;
 (g) $Z(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
 (h) $Z(f, g, \cdot)$ is non decreasing for every value $f \in Q, g \in L$;
 (i) $K(f, g, 0) < 1$, for every value $f, g \in Q \times L$;
 (j) $K(f, g, \Theta) = 1$ iff $f = l$ for $f \in Q, g \in L$;
 (k) $K(f, g, \Theta) = K(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
 (l) $K(f_1, g_2, \Theta + x + z) \geq K(f_1, g_1, \Theta) \odot K(f_2, g_1, x) \odot K(f_2, g_2, z)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L \forall \Theta, x, z > 0$;
 (m) $K(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
 (n) $K(f, g, \cdot)$ is non increasing for every value $f \in Q, g \in L$;
 (o) $Y(f, g, 0) < 1$, for every value $f, g \in Q \times L$;
 (p) $Y(f, g, \Theta) = 1$ iff $f = l$ for $f \in Q, g \in L$;
 (q) $Y(f, g, \Theta) = Y(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
 (r) $Y(f_1, g_2, \Theta + x + z) \geq Y(f_1, g_1, \Theta) \odot D(f_2, g_1, x) \odot D(f_2, g_2, z)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L \forall \Theta, x, z > 0$;
 (s) $Y(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
 (t) $Y(f, g, \cdot)$ is non increasing for every value $f \in Q, g \in L$.

Then, $(Q, L, Z, K, Y, \otimes, \odot)$ is called a *NBMS*.

Definition 2.5. [19] Suppose $Q, L \neq \emptyset$ and $\Upsilon : Q \times L \rightarrow [1, +\infty)$ are considered as an incompetant mappings, \otimes as *CTN* and \odot as *CTC* and Z, K, Y are *NS* on $Q \times L \times (0, +\infty)$ is characterized *NCBMS*, if for each one $(Q, L, Z, K, Y, \otimes, \odot)$ fulfills all $f \in Q, g \in L$ and $x, \Theta, z > 0$ holds the following:

- (a) $0 \leq Z(f, g, \Theta) \leq 1; 0 \leq K(f, g, \Theta) \leq 1; 0 \leq Y(f, g, \Theta) \leq 1$, for every value $(f, g) \in Q \times L$;
 (b) $Z(f, g, \Theta) + K(f, g, \Theta) + Y(f, g, \Theta) \leq 3$;
 (c) $Z(f, g, 0) = 0$, for every value $(f, g) \in Q \times L$;
 (d) $Z(f, g, \Theta) = 1$ iff $f = g$ for $f \in Q, g \in L$;
 (e) $Z(f, g, \Theta) = Z(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
 (f) $Z(f_1, g_2, \Theta + x + z) \geq Z\left(f_1, g_1, \frac{\Theta}{\Upsilon(f_1, l_1)}\right) \otimes Z\left(f_2, g_1, \frac{x}{\Upsilon(f_2, l_1)}\right) \otimes Z\left(f_2, g_2, \frac{z}{\Upsilon(f_2, l_2)}\right)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L$;
 (g) $Z(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
 (h) $Z(f, g, \cdot)$ is non decreasing for every value $f \in Q, g \in L$;
 (i) $K(f, g, 0) = 1$;
 (j) $K(f, g, \Theta) = 0$ iff $f = g$ for $f \in Q, g \in L$;
 (k) $K(f, g, \Theta) = K(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
 (l) $K(f_1, g_2, \Theta + x + z) \leq K\left(f_1, g_1, \frac{\Theta}{\Upsilon(f_1, l_1)}\right) \otimes K\left(f_2, g_1, \frac{x}{\Upsilon(f_2, l_1)}\right) \otimes K\left(f_2, g_2, \frac{z}{\Upsilon(f_2, l_2)}\right)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L$;

- (m) $K(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
- (n) $K(f, g, \cdot)$ is non increasing for every value $f \in Q, g \in L$;
- (o) $Y(f, g, 0) = 1$;
- (p) $Y(f, g, \Theta) = 0$ iff $f = g$ for $f \in Q, g \in L$;
- (q) $Y(f, g, \Theta) = Y(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
- (r) $Y(f_1, g_2, \Theta + x + z) \leq Y\left(f_1, g_1, \frac{\Theta}{\Upsilon(f_1, g_1)}\right) \odot D\left(f_2, g_1, \frac{x}{\Upsilon(f_2, g_1)}\right) \odot D\left(f_2, g_2, \frac{z}{\Upsilon(f_2, g_2)}\right)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L$;
- (s) $Y(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
- (t) $Y(f, g, \cdot)$ is non increasing for every value $f \in Q, g \in L$.
- Then, $(Q, L, Z, K, Y, \otimes, \odot)$ is called a *NCBMS*.

Example 2.6. Let $Q = [0, 1], L = [1, \infty)$. Define Z, K, D are *NS* on $Q \times Q \times (0, +\infty)$ as

$$Z(f, g, \Theta) = \frac{\Theta}{\Theta + d(f, g)}, K(f, g, \Theta) = \frac{d(f, g)}{\Theta + d(f, g)}, Y(f, l, \Theta) = \frac{d(f, g)}{\Theta}$$

with the *CTN* \otimes such that $\Theta_1 \otimes \Theta_2 = \min\{\Theta_1, \Theta_2\}$ and \odot as *CTC* defined as $\Theta_1 \odot \Theta_2 = \max\{\Theta_1, \Theta_2\}$. Define $\Upsilon : Q \times Q \rightarrow [1, +\infty)$ as

$$\Upsilon(f, g) = \begin{cases} 1 & \text{if } f \in Q \text{ and } g \in L \\ \max\{f, g\} & \text{otherwise} \end{cases}$$

Then $(Q, L, Z, K, Y, \otimes, \odot)$ be a *NCBMS*.

Definition 2.7. [16] Suppose $Q, L \neq \emptyset$ and $\Upsilon, \varrho, \chi : Q \times L \rightarrow [1, +\infty)$ are considered as an incompetent mappings, \otimes as *CTN* and Z are *NS* on $Q \times L \times (0, +\infty)$ is characterized *FTCBMS* on Q , if for each one (Q, L, Z, \otimes) fulfills all $f \in Q, g \in L$ and $x, \Theta, z > 0$ holds the following:

- (a) $Z(f, g, \Theta) > 0$, for every value $(f, g) \in Q \times L$;
- (b) $Z(f, g, \Theta) = 1$ iff $f = g$ for $f \in Q, g \in L$;
- (c) $Z(f, g, \Theta) = Z(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
- (d) $Z(f_1, g_2, \Theta + x + z) \geq Z\left(f_1, g_1, \frac{\Theta}{\Upsilon(f_1, g_1)}\right) \otimes Z\left(f_2, g_1, \frac{x}{\Upsilon(f_2, g_1)}\right) \otimes Z\left(f_2, g_2, \frac{z}{\Upsilon(f_2, g_2)}\right)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L$;
- (e) $Z(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS* $Z(f, g, \cdot)$ is non decreasing for every value $f \in Q, g \in L$.
- Then, $(Q, L, Z, K, Y, \otimes, \odot)$ is called a *FTCBMS*.

3. Main results

Definition 3.1. Suppose $Q, L \neq \emptyset$ and $\Upsilon, \varrho, \chi : Q \times L \rightarrow [1, +\infty)$ are considered as a incompetent mappings, \otimes as *CTN* and \odot as *CTC* and Z, K, Y are *NS* on $Q \times L \times (0, +\infty)$ is characterized *NTCBMS*, if for each one $(Q, L, Z, K, Y, \otimes, \odot)$ fulfills all $f \in Q, g \in L$ and $x, \Theta, z > 0$ holds the following:

- (a) $0 \leq Z(f, g, \Theta) \leq 1; 0 \leq K(f, g, \Theta) \leq 1; 0 \leq Y(f, g, \Theta) \leq 1$, for every value $(f, g) \in Q \times L$;
- (b) $Z(f, g, \Theta) + K(f, g, \Theta) + Y(f, g, \Theta) \leq 3$;
- (c) $Z(f, g, \Theta) > 0$, for every value $(f, g) \in Q \times L$;
- (d) $Z(f, g, \Theta) = 1$ iff $f = g$ for $f \in Q, g \in L$;
- (e) $Z(f, g, \Theta) = Z(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
- (f) $Z(f_1, g_2, \Theta + x + z) \geq Z\left(f_1, g_1, \frac{\Theta}{\Upsilon(f_1, g_1)}\right) \otimes Z\left(f_2, g_1, \frac{x}{\Upsilon(f_2, g_1)}\right) \otimes Z\left(f_2, g_2, \frac{z}{\Upsilon(f_2, g_2)}\right)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L$;
- (g) $Z(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
- (h) $Z(f, g, \cdot)$ is non decreasing $\forall f \in Q, g \in L$;
- (i) $K(f, g, \Theta) < 1$;
- (j) $K(f, g, \Theta) = 0$ iff $f = g$ for $f \in Q, g \in L$;
- (k) $K(f, g, \Theta) = K(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
- (l) $K(f_1, g_2, \Theta + x + z) \leq K\left(f_1, g_1, \frac{\Theta}{\Upsilon(f_1, g_1)}\right) \odot K\left(f_2, g_1, \frac{x}{\Upsilon(f_2, g_1)}\right) \odot K\left(f_2, g_2, \frac{z}{\Upsilon(f_2, g_2)}\right)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L$;

- (m) $K(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
 - (n) $K(f, g, \cdot)$ is non increasing for every value $f \in Q, g \in L$;
 - (o) $Y(f, g, \Theta) < 1$;
 - (p) $Y(f, g, \Theta) = 0$ iff $f = g \forall f \in Q, g \in L$;
 - (q) $Y(f, g, \Theta) = Y(g, f, \Theta) \forall f, g \in Q \cap L$;
 - (r) $Y(f_1, g_2, \Theta + x + z) \leq Y\left(f_1, g_1, \frac{\Theta}{\Upsilon(f_1, g_1)}\right) \odot D\left(f_2, g_1, \frac{x}{\Upsilon(f_2, g_1)}\right) \odot D\left(f_2, g_2, \frac{z}{\Upsilon(f_2, g_2)}\right)$ for $f_1, f_2 \in Q$ and $g_1, g_2 \in L$;
 - (s) $Y(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
 - (t) $Y(f, g, \cdot)$ is non increasing $\forall f \in Q, g \in L$;
- Then, $(Q, L, Z, K, Y, \otimes, \odot)$ is called a *NTCBMS*.

Definition 3.2. Suppose $Q, L \neq \emptyset$ and $\Upsilon_i : Q \times L \rightarrow [1, +\infty) (1 \leq i \leq n)$ are considered as a incompetent mappings, \otimes as *CTN* and \odot as *CTC* and Z, K, Y are *NS* on $Q \times L \times (0, +\infty)$ is characterized *NnCBMS*, if for each one $(Q, L, Z, K, Y, \otimes, \odot)$ fulfills all $f \in Q, g \in L$ and $\Theta_i > 0$ holds the following:

- (a) $0 \leq Z(f, g, \Theta) \leq 1; 0 \leq K(f, g, \Theta) \leq 1; 0 \leq Y(f, g, \Theta) \leq 1$, for every value $(f, g) \in Q \times L$;
 - (b) $Z(f, g, \Theta) + K(f, g, \Theta) + Y(f, g, \Theta) \leq 3$;
 - (c) $Z(f, g, \Theta) > 0$, for every value $(f, g) \in Q \times L$;
 - (d) $Z(f, g, \Theta) = 1$ iff $f = g \forall f \in Q, g \in L$;
 - (e) $Z(f, g, \Theta) = Z(g, f, \Theta)$, for every value $f, g \in Q \cap L$;
 - (f) $Z(f_1, g_\theta, \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4 \dots + \Theta_n) \geq Z\left(f_1, g_1, \frac{\Theta_1}{\Upsilon_1(f_1, g_1)}\right) \otimes Z\left(f_2, g_1, \frac{\Theta_2}{\Upsilon_2(f_2, g_1)}\right) \otimes Z\left(f_2, g_2, \frac{\Theta_3}{\Upsilon_3(f_2, g_2)}\right) \dots \otimes Z\left(f_\theta, g_{\theta-1}, \frac{\Theta_{n-1}}{\Upsilon_{n-1}(f_\theta, g_{\theta-1})}\right) \otimes Z\left(f_\theta, g_\theta, \frac{\Theta_n}{\Upsilon_n(f_\theta, g_\theta)}\right) \forall f_1, f_2 \dots f_\theta \in Q$ and $g_1, g_2 \dots g_\theta \in L$;
 - (g) $Z(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
 - (h) $Z(f, g, \cdot)$ is non decreasing for every value $f \in Q, g \in L$;
 - (i) $K(f, g, \Theta) < 1$, for every value $(f, g) \in Q \times L$;
 - (j) $K(f, g, \Theta) = 0$ iff $f = g \forall f \in Q, g \in L$;
 - (k) $K(f, g, \Theta) = K(g, f, \Theta) \forall f, g \in Q \cap L$;
 - (l) $K(f_1, g_\theta, \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4 \dots + \Theta_n) \leq K\left(f_1, g_1, \frac{\Theta_1}{\Upsilon_1(f_1, g_1)}\right) \odot K\left(f_2, g_1, \frac{\Theta_2}{\Upsilon_2(f_2, g_1)}\right) \odot K\left(f_2, g_2, \frac{\Theta_3}{\Upsilon_3(f_2, g_2)}\right) \dots \odot K\left(f_\theta, g_{\theta-1}, \frac{\Theta_{n-1}}{\Upsilon_{n-1}(f_\theta, g_{\theta-1})}\right) \odot K\left(f_\theta, g_\theta, \frac{\Theta_n}{\Upsilon_n(f_\theta, g_\theta)}\right) \forall f_1, f_2 \dots f_\theta \in Q$ and $g_1, g_2 \dots g_\theta \in L$;
 - (m) $K(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
 - (n) $K(f, g, \cdot)$ is non increasing for every value $f \in Q, g \in L$;
 - (o) $Y(f, g, \Theta) < 1 \forall (f, g) \in Q \times L$;
 - (p) $Y(f, g, \Theta) = 0$ iff $f = g \forall f \in Q, g \in L$;
 - (q) $Y(f, g, \Theta) = Y(g, f, \Theta) \forall f, g \in Q \cap L$;
 - (r) $Y(f_1, g_\theta, \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4 \dots + \Theta_n) \leq \left(f_1, g_1, \frac{\Theta_1}{\Upsilon_1(f_1, g_1)}\right) \odot \left(f_2, g_1, \frac{\Theta_2}{\Upsilon_2(f_2, g_1)}\right) \odot Y\left(f_2, g_2, \frac{\Theta_3}{\Upsilon_3(f_2, g_2)}\right) \dots \odot D\left(f_\theta, g_{\theta-1}, \frac{\Theta_{n-1}}{\Upsilon_{n-1}(f_\theta, g_{\theta-1})}\right) \odot Y\left(f_\theta, g_\theta, \frac{\Theta_n}{\Upsilon_n(f_\theta, g_\theta)}\right)$ for $f_1, f_2 \dots f_\theta \in Q$ and $g_1, g_2 \dots g_\theta \in L$;
 - (s) $Y(f, g, \cdot) : [0, +\infty) \rightarrow [0, 1]$ is *CTS*;
 - (t) $Y(f, g, \cdot)$ is non increasing for every value $f \in Q, g \in L$;
- Then, $(Q, L, Z, K, Y, \otimes, \odot)$ is called a *NnCBMS*.

Example 3.3. Let $Q = [0, \frac{1}{2^n}] , L = [\frac{1}{2^{n+1}}, 1]$ and $\Upsilon_i : Q \times L \rightarrow [1, \infty), (1 \leq i \leq 9)$ be nine non-comparable mappings defined as

$$\begin{aligned} \Upsilon_1(f_1, g_1) &= f_1 + g_1 + 1, \Upsilon_2(f_2, g_1) = f_2 + g_1 + 1, \Upsilon_3(f_2, g_2) = f_2 + g_2 + 1, \\ \Upsilon_4(f_3, g_2) &= f_3 + g_2 + 1, \Upsilon_5(f_3, g_3) = f_3 + g_3 + 1, \Upsilon_6(f_4, g_3) = f_4 + g_3 + 1, \\ \Upsilon_7(f_4, g_4) &= f_4 + g_4 + 1, \Upsilon_8(f_5, g_4) = f_5 + g_4 + 1, \Upsilon_9(f_5, g_5) = f_5 + g_5 + 1. \end{aligned}$$

Define $Z(f_1, g_5, \Theta_i) = \frac{\min(f, g) + \Theta}{\max(f, g) + \Theta}$, for every value $\Theta_i > 0, K(f_1, g_5, \Theta_i) = \frac{\max(f, g) - \min(f, g)}{\max(f, g) + \Theta}$, for every value

$\Theta_i > 0$ and $Y(f_1, g_5, \Theta_i) = \frac{\max(f, g) - \min(f, g)}{\min(f, g) + \Theta}$, for every value $\Theta_i > 0, f \in Q$ and $g \in L$. Clearly, $(Q, L, Z, K, Y, \otimes, \odot)$ is a *NnCBMS*, where \otimes is a *CTC* defined as $\Theta_1 \otimes \Theta_2 = \Theta_1 \Theta_2$ and \odot is a *CTN* defined as $\Theta_1 \odot \Theta_2 = \max\{\Theta_1, \Theta_2\}$.

Proof

The first three conditions are met easily so we go for fourth condition *Z, K, Y* as follows:

For $f_i \neq g_i$ and $\Theta_i \geq 0$.

By assuming $f_1 = 0, f_2 = \frac{1}{6}, f_3 = \frac{1}{5}, f_4 = \frac{2}{5}, f_5 = \frac{1}{2}, g_1 = \frac{1}{4}, g_2 = \frac{2}{5}, g_3 = \frac{1}{2}, g_4 = \frac{3}{4}, g_5 = 1$ we obtain a non-trivial sequence as

$$(f_1, g_5) = \{(f_1, g_1), (f_2, g_1), (f_2, g_2), (f_3, g_2), (f_3, g_3), (f_4, g_3), (f_4, g_4), (f_5, g_4), (f_5, g_5)\},$$

$$(0, 1) = \{(0, \frac{1}{4}), (\frac{1}{6}, \frac{1}{4}), (\frac{1}{6}, \frac{2}{5}), (\frac{1}{5}, \frac{2}{5}), (\frac{1}{5}, \frac{1}{2}), (\frac{2}{5}, \frac{1}{2}), (\frac{2}{5}, \frac{3}{4}), (\frac{1}{2}, \frac{3}{4}), (\frac{1}{2}, 1)\}$$
 and taking $\Theta_i = 1$.

$$Z(f_1, g_5, \Theta_i) = \frac{\min(f, g) + \Theta}{\max(f, g) + \Theta}, \text{ for every value } \Theta_i > 0,$$

$$\begin{aligned} Z(0, 1, 9) &\geq Z\left(0, \frac{1}{4}, \frac{1}{\Upsilon_1(0, \frac{1}{4})}\right) \otimes Z\left(\frac{1}{6}, \frac{1}{4}, \frac{1}{\Upsilon_2(\frac{1}{6}, \frac{1}{4})}\right) \otimes Z\left(\frac{1}{6}, \frac{2}{5}, \frac{1}{\Upsilon_3(\frac{1}{6}, \frac{2}{5})}\right) \otimes Z\left(\frac{1}{5}, \frac{2}{5}, \frac{1}{\Upsilon_4(\frac{1}{5}, \frac{2}{5})}\right) \\ &\otimes Z\left(\frac{1}{5}, \frac{1}{2}, \frac{1}{\Upsilon_5(\frac{1}{5}, \frac{1}{2})}\right) \otimes Z\left(\frac{2}{5}, \frac{1}{2}, \frac{1}{\Upsilon_6(\frac{2}{5}, \frac{1}{2})}\right) \otimes Z\left(\frac{2}{5}, \frac{3}{4}, \frac{1}{\Upsilon_7(\frac{2}{5}, \frac{3}{4})}\right) \otimes \\ &Z\left(\frac{1}{2}, \frac{3}{4}, \frac{1}{\Upsilon_8(\frac{1}{2}, \frac{3}{4})}\right) \otimes Z\left(\frac{1}{2}, 1, \frac{1}{\Upsilon_9(\frac{1}{2}, 1)}\right) \\ &= Z\left(f_1, g_1, \frac{\Theta_1}{\Upsilon_1(f_1, l_1)}\right) \otimes Z\left(f_2, g_1, \frac{\Theta_2}{\Upsilon_2(f_2, l_1)}\right) \otimes Z\left(f_2, g_2, \frac{\Theta_3}{\Upsilon_3(f_2, l_2)}\right) \dots \otimes \\ &Z\left(f_\theta, g_{\theta-1}, \frac{\Theta_{n-1}}{\Upsilon_{n-1}(f_\theta, l_{\theta-1})}\right) \otimes Z\left(f_\theta, g_\theta, \frac{\Theta_n}{\Upsilon_n(f_\theta, l_\theta)}\right) \end{aligned}$$

$$K(f_1, g_5, \Theta_i) = \frac{\max(f, l) - \min(f, l)}{\max(f, l) + \Theta}, \text{ for every value } \Theta_i > 0$$

$$\begin{aligned} K(0, 1, 9) &\leq K\left(0, \frac{1}{4}, \frac{1}{\Upsilon_1(0, \frac{1}{4})}\right) \odot K\left(\frac{1}{6}, \frac{1}{4}, \frac{1}{\Upsilon_2(\frac{1}{6}, \frac{1}{4})}\right) \odot K\left(\frac{1}{6}, \frac{2}{5}, \frac{1}{\Upsilon_3(\frac{1}{6}, \frac{2}{5})}\right) \odot K\left(\frac{1}{5}, \frac{2}{5}, \frac{1}{\Upsilon_4(\frac{1}{5}, \frac{2}{5})}\right) \\ &\odot K\left(\frac{1}{5}, \frac{1}{2}, \frac{1}{\Upsilon_5(\frac{1}{5}, \frac{1}{2})}\right) \odot K\left(\frac{2}{5}, \frac{1}{2}, \frac{1}{\Upsilon_6(\frac{2}{5}, \frac{1}{2})}\right) \\ &= K\left(f_1, g_1, \frac{\Theta_1}{\Upsilon_1(f_1, g_1)}\right) \odot K\left(f_2, g_1, \frac{\Theta_2}{\Upsilon_2(f_2, g_1)}\right) \odot K\left(f_2, g_2, \frac{\Theta_3}{\Upsilon_3(f_2, g_2)}\right) \dots \odot \\ &K\left(f_\theta, g_{\theta-1}, \frac{\Theta_{n-1}}{\Upsilon_{n-1}(f_\theta, g_{\theta-1})}\right) \odot K\left(f_\theta, g_\theta, \frac{\Theta_n}{\Upsilon_n(f_\theta, g_\theta)}\right) \end{aligned}$$

$$Y(f_1, g_5, \Theta_i) = \frac{\max(f, l) - \min(f, l)}{\max(f, l) + \Theta}, \text{ for every value } \Theta_i > 0$$

$$\begin{aligned} Y(0, 1, 9) &\leq Y\left(0, \frac{1}{4}, \frac{1}{\Upsilon_1(0, \frac{1}{4})}\right) \odot Y\left(\frac{1}{6}, \frac{1}{4}, \frac{1}{\Upsilon_2(\frac{1}{6}, \frac{1}{4})}\right) \odot Y\left(\frac{1}{6}, \frac{2}{5}, \frac{1}{\Upsilon_3(\frac{1}{6}, \frac{2}{5})}\right) \odot \\ &Y\left(\frac{1}{5}, \frac{2}{5}, \frac{1}{\Upsilon_4(\frac{1}{5}, \frac{2}{5})}\right) \odot Y\left(\frac{1}{5}, \frac{1}{2}, \frac{1}{\Upsilon_5(\frac{1}{5}, \frac{1}{2})}\right) \odot Y\left(\frac{2}{5}, \frac{1}{2}, \frac{1}{\Upsilon_6(\frac{2}{5}, \frac{1}{2})}\right) \\ &= Y\left(f_1, g_1, \frac{\Theta_1}{\Upsilon_1(f_1, g_1)}\right) \odot Y\left(f_2, g_1, \frac{\Theta_2}{\Upsilon_2(f_2, g_1)}\right) \odot Y\left(f_2, g_2, \frac{\Theta_3}{\Upsilon_3(f_2, g_2)}\right) \dots \odot \\ &Y\left(f_\theta, g_{\theta-1}, \frac{\Theta_{n-1}}{\Upsilon_{n-1}(f_\theta, g_{\theta-1})}\right) \odot Y\left(f_\theta, g_\theta, \frac{\Theta_n}{\Upsilon_n(f_\theta, g_\theta)}\right), \text{ for every value } \Theta_i > 0. \end{aligned}$$

Similarly we can prove other cases.

The following diagram shows the graphical behaviour of the above example: By calculating the above, which meets




Figure 1. Shows the graphical behaviour of Z, K and Y in the above example

the necessary criterion. We are also able to prove other cases. Similarly, we can demonstrate a greater value of n . Thus, $(Q, L, Z, K, Y, \otimes, \odot)$ is a $NnCBMS$. □

Lemma 3.4. Let $(Q, L, Z, K, Y, \otimes, \odot)$ be a $NnCBMS$ given by $Z(f, g, \delta\Theta) \geq Z(f, g, \Theta)$ for, $f \in Q, g \in L, \Theta \in (0, +\infty)$ and $\delta \in (0, 1)$. Then $f = g$.

Proof

Given that,

$$Z(f, g, \delta\Theta) \geq Z(f, g, \Theta), K(f, g, \delta\Theta) \leq K(f, g, \Theta), Y(f, g, \delta\Theta) \leq Y(f, g, \Theta) \text{ for } \Theta > 0. \tag{3.4.1}$$

Since $\delta\Theta < \Theta$ for every value $\Theta > 0$ and $\delta \in (0, 1)$, by (h),(n),(t) of definition (3.2), we have

$$Z(f, g, \delta\Theta) \geq Z(f, g, \Theta), K(f, g, \delta\Theta) \leq K(f, g, \Theta), Y(f, g, \delta\Theta) \leq Y(f, g, \Theta) \text{ for } \Theta > 0 \tag{3.4.2}$$

From the definition of $NnCBMS$ (see (d),(j),(p)), we obtain $f = g$. □

Definition 3.5. If every Cauchy bisequence is present, it is considered as complete then $(Q, L, Z, K, Y, \otimes, \odot)$ be a $NnCBMS$ in $Q \times L$ is convergent in it.

Proposition 3.6. In a $NnCBMS$, every convergent has a biconvergence in the Cauchy bisequence .

Proof

Let $(Q, L, Z, K, Y, \otimes, \odot)$ be a $NnCBMS$ and a biconvergence $(f_n, g_n) \in Q \times L$ such that $f_n \rightarrow g$ as $n \rightarrow +\infty$ and $g_n \rightarrow f$ as $n \rightarrow +\infty$, where $n \in L$ and $f \in Q$. Since (f_n, g_n) is a convergent Cauchy bisequence , we obtain

$$Z(f_n, g_n, \Theta) \rightarrow 1, K(f_n, g_n, \Theta) \rightarrow 0 \text{ and } Y(f_n, g_n, \Theta) \rightarrow 0 \text{ as } n \rightarrow \infty \tag{3.6.1}$$

for every value $\Theta > 0$. Now, from (3), we conclude that $Z(f, g, \Theta) = 1, K(f, g, \Theta) = 0$ and $Y(f, g, \Theta) = 0$ for every value $\Theta > 0$. Therefore, by (f),(g),(r) of definition (3.2) we obtain that the biconvergence (f_n, g_n) is biconvergent □

Proposition 3.7. In a $NnCBMS$, every biconvergent bisequence is a Cauchy bisequence.

Proof

Let $(Q, L, Z, K, Y, \otimes, \odot)$ be a $NnCBMS$ and a biconvergence $(f_n, g_n) \in Q \times L$ converges to a point $f_0 \in Q \cap L$ $\forall n, \theta \in \mathbb{N}$ and $\Theta, e > 0$. By (f),(g),(r) of definition (3.2), we have biconvergence

$$\begin{aligned} Z(f_n, g_\theta, \Theta) &\geq Z\left(f_n, f_0, \frac{\Theta}{\Upsilon_1(f_n, f_0)}\right) \otimes Z\left(f_0, f_0, \frac{\Theta}{\Upsilon_2(f_0, f_0)}\right) \otimes Z\left(f_0, f_0, \frac{\Theta}{\Upsilon_3(f_0, f_0)}\right) \\ &\quad \otimes Z\left(f_0, f_0, \frac{\Theta}{\Upsilon_4(f_0, f_0)}\right) \cdots \otimes Z\left(f_0, g_\omega, \frac{\Theta}{\Upsilon_n(\vartheta_0, \vartheta_\theta)}\right), \\ K(f_n, g_\theta, \Theta) &\leq K\left(f_n, f_0, \frac{\Theta}{\Upsilon_1(f_n, f_0)}\right) \odot K\left(f_0, f_0, \frac{\Theta}{\Upsilon_2(f_0, f_0)}\right) \odot K\left(f_0, f_0, \frac{\Theta}{\Upsilon_3(f_0, f_0)}\right) \\ &\quad \odot K\left(f_0, f_0, \frac{\Theta}{\Upsilon_4(f_0, f_0)}\right) \cdots \odot K\left(f_0, g_\omega, \frac{\Theta}{\Upsilon_n(\vartheta_0, \vartheta_\theta)}\right), \\ Y(f_n, g_\theta, \Theta) &\leq Y\left(f_n, f_0, \frac{\Theta}{\Upsilon_1(f_n, f_0)}\right) \odot Y\left(f_0, f_0, \frac{\Theta}{\Upsilon_2(f_0, f_0)}\right) \odot Y\left(f_0, f_0, \frac{\Theta}{\Upsilon_3(f_0, f_0)}\right) \\ &\quad \odot Y\left(f_0, f_0, \frac{\Theta}{\Upsilon_4(f_0, f_0)}\right) \cdots \odot Y\left(f_0, g_\omega, \frac{\Theta}{\Upsilon_n(\vartheta_0, \vartheta_\theta)}\right). \end{aligned}$$

As $n, \theta \rightarrow \infty$, we obtain $Z(f_n, g_\theta, \Theta) \geq 1, K(f_n, g_\theta, \Theta) \leq 0$ and $Y(f_n, g_\theta, \Theta) \leq 0$ for every value $\Theta, e > 0$. This implies that $Z(f_n, g_\theta, \Theta) \rightarrow 1, K(f_n, g_\theta, \Theta) \rightarrow 0$ and $Y(f_n, g_\theta, \Theta) \rightarrow 0$ for every value $\Theta, e > 0$. Hence (f_n, g_n) is a Cauchy bisequence. \square

Lemma 3.8. Let $(Q, L, Z, K, Y, \otimes, \odot)$ be a $NnCBMS$ and $\Omega \in Q \cap L$ is a sequence's limit, then it is the sequence's unique limit.

Proof

Get $\{f_n\} \in Q$ be a sequence. Consider that $\{f_n\} \rightarrow g \in L$ and also $\{f_n\} \rightarrow \Omega \in Q \cap L$, then for $\Theta_i > 0$, we have

$$\begin{aligned} Z(\Omega, g, \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4 \cdots + \Theta_n) &\geq Z(\Omega, \Omega, \Theta_1) \otimes Z(\Omega, \Omega, \Theta_2) \otimes Z(\Omega, \Omega, \Theta_3) \otimes \cdots \otimes Z(f_n, \Omega, \Theta_{n-1}) \otimes Z(f_n, g, \Theta_n), \\ K(\Omega, g, \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4 \cdots + \Theta_n) &\leq K(\omega, \omega, \Theta_1) \odot K(\omega, \omega, \Theta_2) \odot K(\omega, \omega, \Theta_3) \odot \cdots \odot K(f_n, \omega, \Theta_{n-1}) \odot K(f_n, g, \Theta_n), \\ Y(\Omega, g, \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4 \cdots + \Theta_n) &\leq Y(\Omega, \Omega, \Theta_1) \odot Y(\Omega, \Omega, \Theta_2) \odot Y(\Omega, \Omega, \Theta_3) \odot \cdots \odot Y(f_n, \Omega, \Theta_{n-1}) \odot Y(f_n, g, \Theta_n). \end{aligned}$$

As $n \rightarrow \infty$, we obtain $Z(\Omega, g, \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4 \cdots + \Theta_n) \geq 1, K(\Omega, g, \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4 \cdots + \Theta_n) \leq 0, Y(\Omega, g, \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4 \cdots + \Theta_n) \leq 0$ which suggest that $\Omega = g$, i.e., sequence $\{f_n\}$, has a unique limit. \square

Theorem 3.9. Let $(Q, L, Z, K, Y, \otimes, \odot)$ be a complete $NnCBMS$ with n non-comparable functions $\Upsilon_i : Q \times L \rightarrow [1, \infty)$ given by

$$\lim_{\Theta \rightarrow \infty} Z(f, g, \Theta) = 1, \lim_{\Theta \rightarrow \infty} K(f, g, \Theta) = 0, \lim_{\Theta \rightarrow \infty} Y(f, g, \Theta) = 0, \text{ for every value } f \in Q, g \in L. \tag{3.9.1}$$

Get $\Xi : Q \cup L \rightarrow Q \cup L$ be a mapping satisfying

- (i) $\Xi(Q) \subseteq Q$ and $\Xi(L) \subseteq L$;
- (ii) $\prod_{\rho} (\Xi(f), \Xi(g), \delta\Theta) \geq Z(f, g, \Theta)$ for every value $f \in Q, g \in L$, and $\Theta > 0$, where $\delta \in (0, 1)$. Assume moreover that $\lim_{\eta \rightarrow \infty} \Upsilon_i(f_n, g)$ and $\lim_{\eta \rightarrow \infty} \Upsilon_i(g, f_n)$, exist and are finite for each $f \in Q$. Then Ξ has a *FP* that is unique.

Proof

Fix $f_0 \in Q$ and $g_0 \in L$ and consider that $\Xi(f_n) = f_{n+1}$ and $\Xi(g_n) = g_{n+1} \forall n \in \mathbb{N} \cup \{0\}$. Next, we get (f_n, g_n) as a biconvergence on $NnCBMS (Q, L, Z, K, Y, \otimes, \odot)$. Now, we have

$$\begin{aligned} Z(f_1, g_1, \Theta) &= Z(\Xi(f_0), \Xi(g_0), \Theta) \geq Z\left(f_0, g_0, \frac{\Theta}{\delta}\right), \\ K(f_1, g_1, \Theta) &= K(\Xi(f_0), \Xi(g_0), \Theta) \leq K\left(f_0, g_0, \frac{\Theta}{\delta}\right), \\ Y(f_1, g_1, \Theta) &= Y(\Xi(f_0), \Xi(g_0), \Theta) \leq Y\left(f_0, g_0, \frac{\Theta}{\delta}\right) \forall \Theta > 0 \text{ and } n \in \mathbb{N}. \end{aligned}$$

By simple induction, we get,

$$\begin{aligned} Z(f_n, g_n, \Theta) &= Z(\Xi(f_{n-1}), \Xi(g_{n-1}), \Theta) \geq Z\left(f_0, g_0, \frac{\Theta^n}{\delta}\right), \\ K(f_n, g_n, \Theta) &= K(\Xi(f_{n-1}), \Xi(g_{n-1}), \Theta) \leq K\left(f_0, g_0, \frac{\Theta^n}{\delta}\right), \\ Y(f_n, g_n, \Theta) &= Y(\Xi(f_{n-1}), \Xi(g_{n-1}), \Theta) \leq Y\left(f_0, g_0, \frac{\Theta^n}{\delta}\right), \end{aligned} \tag{3.9.2}$$

and

$$\begin{aligned} Z(f_{n+1}, g_n, \Theta) &= Z(\Xi(f_n), \Xi(g_{n-1}), \Theta) \geq Z\left(f_1, g_0, \frac{\Theta^n}{\delta}\right), \\ K(f_{n+1}, g_n, \Theta) &= K(\Xi(f_n), \Xi(g_{n-1}), \Theta) \leq K\left(f_1, g_0, \frac{\Theta^n}{\delta}\right), \\ Y(f_{n+1}, g_n, \Theta) &= Y(\Xi(f_n), \Xi(g_{n-1}), \Theta) \leq Y\left(f_1, g_0, \frac{\Theta^n}{\delta}\right) \forall \Theta > 0 \text{ and } n \in \mathbb{N}. \end{aligned} \tag{3.9.3}$$

Let $n < \omega \forall n, \omega \in \mathbb{N}$. Then

$$\begin{aligned} Z(f_n, g_\omega, \Theta) &\geq Z\left(f_n, g_n, \frac{\Theta}{\Upsilon_1(f_n, g_n)}\right) \otimes Z\left(f_{n+1}, g_n, \frac{\Theta}{\Upsilon_2(f_{n+1}, g_n)}\right) \otimes Z\left(f_{n+1}, g_{n+1}, \frac{\Theta}{\Upsilon_3(f_{n+1}, g_{n+1})}\right) \otimes \\ &Z\left(f_{n+2}, g_{n+1}, \frac{\Theta}{\Upsilon_4(f_{n+2}, g_{n+1})}\right) \otimes \dots \otimes Z\left(f_\omega, g_{\omega-1}, \frac{\Theta}{\Upsilon_{n-1}(f_\omega, g_{\omega-1})}\right) \otimes Z\left(f_\omega, g_\omega, \frac{\Theta}{\Upsilon_n(f_\omega, g_\omega)}\right), \\ K(f_n, g_\omega, \Theta) &\leq K\left(f_n, g_n, \frac{\Theta}{\Upsilon_1(f_n, g_n)}\right) \odot K\left(f_{n+1}, g_n, \frac{\Theta}{\Upsilon_2(f_{n+1}, g_n)}\right) \odot K\left(f_{n+1}, g_{n+1}, \frac{\Theta}{\Upsilon_3(f_{n+1}, g_{n+1})}\right) \odot \\ &K\left(f_{n+2}, g_{n+1}, \frac{\Theta}{\Upsilon_4(f_{n+2}, g_{n+1})}\right) \odot \dots \odot K\left(f_\omega, g_{\omega-1}, \frac{\Theta}{\Upsilon_{n-1}(f_\omega, g_{\omega-1})}\right) \odot K\left(f_\omega, g_\omega, \frac{\Theta}{\Upsilon_n(f_\omega, g_\omega)}\right), \\ Y(f_n, g_\omega, \Theta) &\leq Y\left(f_n, g_n, \frac{\Theta}{\Upsilon_1(f_n, g_n)}\right) \odot Y\left(f_{n+1}, g_n, \frac{\Theta}{\Upsilon_2(f_{n+1}, g_n)}\right) \odot Y\left(f_{n+1}, g_{n+1}, \frac{\Theta}{\Upsilon_3(f_{n+1}, g_{n+1})}\right) \odot \\ &Y\left(f_{n+2}, g_{n+1}, \frac{\Theta}{\Upsilon_4(f_{n+2}, g_{n+1})}\right) \odot \dots \odot Y\left(f_\omega, g_{\omega-1}, \frac{\Theta}{\Upsilon_{n-1}(f_\omega, g_{\omega-1})}\right) \odot Y\left(f_\omega, g_\omega, \frac{\Theta}{\Upsilon_n(f_\omega, g_\omega)}\right). \end{aligned}$$

With (3.9.2) and (3.9.3) applied to each and every term on the RHS of the inequality above, we now get

$$\begin{aligned}
 Z(\mathbb{f}_n, \mathfrak{g}_\omega, \Theta) &\geq Z\left(\mathbb{f}_0, \mathfrak{g}_0, \frac{\Theta}{q^n \Upsilon_1(\mathbb{f}_n, \mathfrak{g}_n)}\right) \otimes Z\left(\mathbb{f}_1, \mathfrak{g}_0, \frac{\Theta}{q^{n+1} \Upsilon_2(\mathbb{f}_{n+1}, \mathfrak{g}_n)}\right) \otimes Z\left(\mathbb{f}_1, \mathfrak{g}_1, \frac{\Theta}{q^{n+2} \Upsilon_3(\mathbb{f}_{n+1}, \mathfrak{g}_{n+1})}\right) \otimes \\
 &Z\left(\mathbb{f}_2, \mathfrak{g}_1, \frac{\Theta}{q^{n+3} \Upsilon_4(\mathbb{f}_{n+2}, \mathfrak{g}_{n+1})}\right) \otimes \cdots \otimes Z\left(\mathbb{f}_1, \mathfrak{g}_0, \frac{\Theta}{q^{\omega-1} \Upsilon_{n-1}(\mathbb{f}_\omega, \mathfrak{g}_{\omega-1})}\right) \otimes Z\left(\mathbb{f}_0, \mathfrak{g}_0, \frac{\Theta}{q^\omega \Upsilon_n(\mathbb{f}_\omega, \mathfrak{g}_\omega)}\right), \\
 K(\mathbb{f}_n, \mathfrak{g}_\omega, \Theta) &\leq K\left(\mathbb{f}_0, \mathfrak{g}_0, \frac{\Theta}{q^n \Upsilon_1(\mathbb{f}_n, \mathfrak{g}_n)}\right) \odot K\left(\mathbb{f}_1, \mathfrak{g}_0, \frac{\Theta}{q^{n+1} \Upsilon_2(\mathbb{f}_{n+1}, \mathfrak{g}_n)}\right) \odot K\left(\mathbb{f}_1, \mathfrak{g}_1, \frac{\Theta}{q^{n+2} \Upsilon_3(\mathbb{f}_{n+1}, \mathfrak{g}_{n+1})}\right) \odot \\
 &K\left(\mathbb{f}_2, \mathfrak{g}_1, \frac{\Theta}{q^{n+3} \Upsilon_4(\mathbb{f}_{n+2}, \mathfrak{g}_{n+1})}\right) \odot \cdots \odot K\left(\mathbb{f}_1, \mathfrak{g}_0, \frac{\Theta}{q^{\omega-1} \Upsilon_{n-1}(\mathbb{f}_\omega, \mathfrak{g}_{\omega-1})}\right) \odot K\left(\mathbb{f}_0, \mathfrak{g}_0, \frac{\Theta}{q^\omega \Upsilon_n(\mathbb{f}_\omega, \mathfrak{g}_\omega)}\right), \\
 Y(\mathbb{f}_n, \mathfrak{g}_\omega, \Theta) &\leq Y\left(\mathbb{f}_0, \mathfrak{g}_0, \frac{\Theta}{q^n \Upsilon_1(\mathbb{f}_n, \mathfrak{g}_n)}\right) \odot Y\left(\mathbb{f}_1, \mathfrak{g}_0, \frac{\Theta}{q^{n+1} \Upsilon_2(\mathbb{f}_{n+1}, \mathfrak{g}_n)}\right) \odot Y\left(\mathbb{f}_1, \mathfrak{g}_1, \frac{\Theta}{q^{n+2} \Upsilon_3(\mathbb{f}_{n+1}, \mathfrak{g}_{n+1})}\right) \odot \\
 &Y\left(\mathbb{f}_2, \mathfrak{g}_1, \frac{\Theta}{q^{n+3} \Upsilon_4(\mathbb{f}_{n+2}, \mathfrak{g}_{n+1})}\right) \odot \cdots \odot Y\left(\mathbb{f}_1, \mathfrak{g}_0, \frac{\Theta}{q^{\omega-1} \Upsilon_{n-1}(\mathbb{f}_\omega, \mathfrak{g}_{\omega-1})}\right) \odot Y\left(\mathbb{f}_0, \mathfrak{g}_0, \frac{\Theta}{q^\omega \Upsilon_n(\mathbb{f}_\omega, \mathfrak{g}_\omega)}\right).
 \end{aligned}$$

From (4), as $n, \omega \rightarrow \infty$, we get $Z(\mathbb{f}_n, \mathfrak{g}_\omega, \Theta) \geq 1, K(\mathbb{f}_n, \mathfrak{g}_\omega, \Theta) \leq 0, Y(\mathbb{f}_n, \mathfrak{g}_\omega, \Theta) \leq 0$ for every value $\Theta > 0$.

Bisqnce $(\mathbb{f}_n, \mathfrak{g}_n)$ is hence a Cauchy bisequence . Given the completeness of $(Q, L, Z, K, Y, \otimes, \odot)$, the biconvergence $(\mathbb{f}_n, \mathfrak{g}_n)$ is a convergent Cauchy bisequence . Proposition (3.6) states that the biconvergence $(\mathbb{f}_n, \mathfrak{g}_n)$ is a biconvergent sequence.

Due to the biconvergence of $(\mathbb{f}_n, \mathfrak{g}_n)$, both the $\{\mathbb{f}_n\}$ and $\{\mathfrak{g}_n\}$ sequences have a limit at $\Omega \in Q \cap L$. Lemma (3.8) states that each of the sequences $\{\mathbb{f}_n\}$ and $\{\mathfrak{g}_n\}$ has a distinct limit. Examine (f), (g), and (r) of definition (3.2).

$$\begin{aligned}
 Z(\Xi(\Omega), \Omega, \Theta) &\geq Z\left(\Xi(\Omega), \Xi(\mathfrak{g}_n), \frac{\Theta}{\Upsilon_1(\Omega, \mathfrak{g}_n)}\right) \otimes Z\left(\Xi(\mathbb{f}_n), \Xi(\mathfrak{g}_n), \frac{\Theta}{\Upsilon_1(\mathbb{f}_n, \mathfrak{g}_n)}\right) \otimes \\
 &Z\left(\Xi(\mathbb{f}_n), \Xi(\mathfrak{g}_{n+1}), \frac{\Theta}{\Upsilon_3(\mathbb{f}_{n+1}, \mathfrak{g}_{n+1})}\right) \otimes Z\left(\Xi(\mathbb{f}_{n+1}), \Xi(\mathfrak{g}_{n+1}), \frac{\Theta}{\Upsilon_4(\mathbb{f}_{n+1}, \mathfrak{g}_{n+1})}\right) \otimes \cdots \otimes \\
 &Z\left(\Xi(\mathbb{f}_{\omega-1}), \Xi(\mathfrak{g}_{\omega-1}), \frac{\Theta}{\Upsilon_{n-1}(\mathbb{f}_{\omega-1}, \mathfrak{g}_{\omega-1})}\right) \otimes Z\left(\Xi(\mathbb{f}_\omega), \Xi(\Omega), \frac{\Theta}{\Upsilon_n(\mathbb{f}_\omega, \Omega)}\right), \\
 K(\Xi(\Omega), \Omega, \Theta) &\leq K\left(\Xi(\Omega), \Xi(\mathfrak{g}_n), \frac{\Theta}{\Upsilon_1(\Omega, \mathfrak{g}_n)}\right) \odot K\left(\Xi(\mathbb{f}_n), \Xi(\mathfrak{g}_n), \frac{\Theta}{\Upsilon_1(\mathbb{f}_n, \mathfrak{g}_n)}\right) \odot \\
 &K\left(\Xi(\mathbb{f}_n), \Xi(\mathfrak{g}_{n+1}), \frac{\Theta}{\Upsilon_3(\mathbb{f}_{n+1}, \mathfrak{g}_{n+1})}\right) \odot K\left(\Xi(\mathbb{f}_{n+1}), \Xi(\mathfrak{g}_{n+1}), \frac{\Theta}{\Upsilon_4(\mathbb{f}_{n+1}, \mathfrak{g}_{n+1})}\right) \odot \cdots \odot \\
 &K\left(\Xi(\mathbb{f}_{\omega-1}), \Xi(\mathfrak{g}_{\omega-1}), \frac{\Theta}{\Upsilon_{n-1}(\mathbb{f}_{\omega-1}, \mathfrak{g}_{\omega-1})}\right) \odot K\left(\Xi(\mathbb{f}_\omega), \Xi(\Omega), \frac{\Theta}{\Upsilon_n(\mathbb{f}_\omega, \Omega)}\right), \\
 Y(\Xi(\Omega), \Omega, \Theta) &\leq Y\left(\Xi(\Omega), \Xi(\mathfrak{g}_n), \frac{\Theta}{\Upsilon_1(\Omega, \mathfrak{g}_n)}\right) \odot Y\left(\Xi(\mathbb{f}_n), \Xi(\mathfrak{g}_n), \frac{\Theta}{\Upsilon_1(\mathbb{f}_n, \mathfrak{g}_n)}\right) \odot \\
 &Y\left(\Xi(\mathbb{f}_n), \Xi(\mathfrak{g}_{n+1}), \frac{\Theta}{\Upsilon_3(\mathbb{f}_{n+1}, \mathfrak{g}_{n+1})}\right) \odot Y\left(\Xi(\mathbb{f}_{n+1}), \Xi(\mathfrak{g}_{n+1}), \frac{\Theta}{\Upsilon_4(\mathbb{f}_{n+1}, \mathfrak{g}_{n+1})}\right) \odot \cdots \odot \\
 &Y\left(\Xi(\mathbb{f}_{\omega-1}), \Xi(\mathfrak{g}_{\omega-1}), \frac{\Theta}{\Upsilon_{n-1}(\mathbb{f}_{\omega-1}, \mathfrak{g}_{\omega-1})}\right) \odot Y\left(\Xi(\mathbb{f}_\omega), \Xi(\Omega), \frac{\Theta}{\Upsilon_n(\mathbb{f}_\omega, \Omega)}\right), \forall n, \omega \in \mathbb{N} \text{ and } \Theta > 0.
 \end{aligned}$$

As $n \rightarrow \infty$, implies

$$\begin{aligned}
 Z(\Xi(\Omega), \Omega, \Theta) &\rightarrow 1 \otimes 1 \otimes 1 \otimes 1 \cdots \otimes 1 \otimes 1 = 1, \\
 K(\Xi(\Omega), \Omega, \Theta) &\rightarrow 0 \odot 0 \odot 0 \odot 0 \cdots \odot 0 \odot 0 = 0,
 \end{aligned}$$

$Y(\Xi(\Omega), \Omega, \Theta) \rightarrow 0 \odot 0 \odot 0 \odot 0 \odot 0 \cdots \odot 0 \odot 0 = 0$.

From (d) of definition (3.2), we obtain $\Xi(\Omega) = \Omega$. Let $\mathbb{f} \in Q \cap L$ be one more *FP* of Ξ . Then

$$\begin{aligned} Z(\Omega, k, \Theta) &= Z(\Xi(\mu), \Xi(\mathbb{f}), \Theta) \geq Z\left(\Omega, k, \frac{\Theta}{q}\right), \\ K(\Omega, k, \Theta) &= K(\Xi(\Omega), \Xi(\mathbb{f}), \Theta) \leq K\left(\Omega, k, \frac{\Theta}{q}\right), \\ Y(\Omega, k, \Theta) &= Y(\Xi(\Omega), \Xi(\mathbb{f}), \Theta) \leq Y\left(\Omega, k, \frac{\Theta}{q}\right). \end{aligned}$$

For $q \in (0, 1)$ and $\forall \Theta > 0$. By Lemma (3.4), finally we get $\Omega = \mathbb{f}$. □

Example 3.10. Let $Q = [0, \frac{1}{2}^n]$, $L = [\frac{1}{2}^{n+1}, 1]$, and $\Upsilon_i : Q \times L \rightarrow [1, \infty)$, ($1 \leq i \leq 9$) be nine non-comparable mappings defined as

$$\Upsilon_1(\mathbb{f}_1, \mathbb{g}_1) = \mathbb{f}_1 + \mathbb{g}_1 + 1, \Upsilon_2(\mathbb{f}_2, \mathbb{g}_1) = \mathbb{f}_2 + \mathbb{g}_1 + 1, \Upsilon_3(\mathbb{f}_2, \mathbb{g}_2) = \mathbb{f}_2 + \mathbb{g}_2 + 1,$$

$$\Upsilon_4(\mathbb{f}_3, \mathbb{g}_2) = \mathbb{f}_3 + \mathbb{g}_2 + 1, \Upsilon_5(\mathbb{f}_3, \mathbb{g}_3) = \mathbb{f}_3 + \mathbb{g}_3 + 1, \Upsilon_6(\mathbb{f}_4, \mathbb{g}_3) = \mathbb{f}_4 + \mathbb{g}_3 + 1,$$

$$\Upsilon_7(\mathbb{f}_4, \mathbb{g}_4) = \mathbb{f}_4 + \mathbb{g}_4 + 1, \Upsilon_8(\mathbb{f}_5, \mathbb{g}_4) = \mathbb{f}_5 + \mathbb{g}_4 + 1, \Upsilon_9(\mathbb{f}_5, \mathbb{g}_5) = \mathbb{f}_5 + \mathbb{g}_5 + 1.$$

Define $Z(\mathbb{f}_1, \mathbb{g}_5, \Theta_i) = \frac{\Theta_i}{\Theta_i + |\mathbb{f}_1 - \mathbb{g}_5|}$, $K(\mathbb{f}_1, \mathbb{g}_5, \Theta_i) = \frac{|\mathbb{f}_1 - \mathbb{g}_5|}{\Theta_i + |\mathbb{f}_1 - \mathbb{g}_5|}$, $Y(\mathbb{f}_1, \mathbb{g}_5, \Theta_i) = \frac{|\mathbb{f}_1 - \mathbb{g}_5|}{\Theta_i}$ for every value $\Theta_i > 0$, $\mathbb{f} \in Q$ and $\mathbb{g} \in L$. Clearly, $(Q, L, Z, K, Y, \otimes, \odot)$ is a *NnCBMS*, where \otimes is a *CTC* defined as $\Theta_1 \otimes \Theta_2 = \Theta_1 \Theta_2$ and \odot is a *CTN* defined as $\Theta_1 \odot \Theta_2 = \max\{\Theta_1, \Theta_2\}$.

Define $\Xi : Q \cup L \rightarrow Q \cup L$ by $\Xi(\Omega) = \begin{cases} \frac{\Omega}{2}, & \text{if } \Omega \in [0, \frac{1}{2}^n], \\ 0, & \text{if } \Omega \in [\frac{1}{2^{n+1}}, 1], \end{cases}$ for every value $\Omega \in Q \cup L$. Obviously, the criteria's of Theorem (3.9) are satisfied. Hence Ξ has a unique *FP*, i.e., $\Omega = 0$. On the same lines, we can prove for higher value of n .

4. Conclusion

We provide the idea of *Nn - CBMS* in this study. Several *FP* outcomes are established using n -non-comparable functions. We also extend the results using a variation of the Banach contraction principle and provide a number of non-trivial instances. We then use the key results to solve fractional differential equation issues related to financial modelling. Future study on additional criteria that would ensure the presence of *FPS* in *Nn - CBMS* is something we find intriguing.

REFERENCES

1. Ali Mutlu, Utku Gurdal Bipolar metric spaces and some fixed point theorems, J. Nonlinear Sci. Appl. 2016, 9(9), 5362-5373.
2. Chen, Haiting, Practical Overview of Triangular Bipolar Neutrosophic Numbers for Design Effect Evaluation of Ethnic Minority Clothing, Comprehensive Guide. Neutrosophic Sets and Systems 81, 1 (2025).
3. George. A, Veeramani. P, On some results in fuzzy metric spaces, Fuzzy Sets Syst. 1994, 64, 395-399.
4. Fahim Uddin, Umar Ishtiaq, Naeem Saleem, Fixed point theorems for controlled neutrosophic metric-like spaces, AIMS Mathematics 2022, vol. 7, 20711-20739.
5. Florentin Smarandache (ed.) Neutrosophic Theory and its Applications, vol.1, Brussels, Europa Nova, 2014.
6. Grabiec. M, Fixed points in fuzzy metric spaces, Fuzzy Sets Syst. 1988, 27, 385-389.
7. Gregori. V, Sapena. A, On fixed point theorems in fuzzy metric spaces, Fuzzy Sets Syst. 2002, 125, 245-252.
8. Gupta. V, Mani. N, Saini. A, Fixed point theorems and its applications in fuzzy metric spaces, Conf. Pap. 2013, AEMDS-2013,961-964.
9. Jeyaraman. M and Shakila. V.B, Common Fixed Point Theorems For A-Admissible Mappings In Neutrosophic Metric Spaces, Advances And Applications In Mathematicag Sciences Volume 21, Issue 4, February 2022, 2069-2082.
10. Jeyaraman. M, Sowndrarajan. S Common Fixed Point Results in Neutrosophic Metric Spaces, Neutrosophic Sets and Systems 42, 1 (2021).
11. Khulud Fahad Bin Muhaya, Kholood Mohammad Alsagar, Extending neutrosophic set theory, Cubic bipolar neutrosophic soft sets for decision making[J]. AIMS Mathematics, 2024,9(10), 27739-27769.

12. Kishore. G.N.V, Agarwag, Ravi. P, Rao. B, Srinivasa. R.V.N, Srinivasa. R, Caristi type contraction and common fixed point theorems in bipolar metric spaces with applications, *Fixed Point Theory Appl.* 2018, 21.
13. Kirisci. M, and Simsek. N, Neutrosophic metric spaces, *Journal of Mathematical Sciences*, 14(28),(2020).
14. Kramosil. I, Michalek. J, Fuzzy metric and statistical metric spaces, *Kybernetika* 1975, 11, 326-334.
15. Mehmood. F, Ali. E, Ionescu. C, Kamran. T, Extended fuzzy b-metric spaces, *J. Math. Anal.* 2017, 8, 124-131.
16. Mitrovic. Z, Bota. M, Solving an Integral Equation via Fuzzy Triple Controlled Bipolar Metric Spaces, *Mathematics* 2021, 9, 3181.
17. Muhiuddin. G, Satham Hussain. S and Durga Nagarajan, Quadri partitioned Bipolar Neutrosophic Competition Graph with Novel Application, *Neutrosophic Sets and Systems* 82, 1 (2025).
18. Rakic. D, Mukheimer. A, Dosenovic. T, Mitrovic. Z.D, Radenovic. S, On some new fixed point results in fuzzy b-metric spaces, *J. Inequal. Appl.* 2020, 99.
19. Rathivel. M, Jeyaraman. M, & Pazhani. V, Investigating Bipolar Controlled Neutrosophic Metric Spaces In Automobile Suspension, *Journal of Computational Analysis and Applications (JoCAAA)*, 33(05), 577-588, (2024).
20. Sezen. M.S, Controlled fuzzy metric spaces and some related fixed point results, *Numer. Methods Partial Differ. Equ.* 2021, 37,583-593.
21. Sowndrarajan. S, Jeyaraman. M and Florentin Smarandache, Fixed Point Results for Contraction Theorems in Neutrosophic Metric Spaces, *Neutrosophic Sets and Systems* 36, 1 (2020).
22. Zadeh. G.A, Fuzzy sets, *Inf. Control* 1965, 8, 338-353.