



# Decision Making Using The Score Function And New Distance Measure: A Comparative Study.

Kamal Abdelraouf Eldahshan<sup>1</sup>, Salah Boulaaras<sup>2,\*</sup>, Ghada E. Mostafa<sup>1</sup>, Zeinab Mohammed<sup>1</sup>,  
Ibrahim Mekawy<sup>3</sup>

<sup>1</sup>Department of Mathematics, Faculty of Science, Al-Azhar University, Cairo, Egypt

<sup>2</sup>Department of Mathematics, College of Sciences, Qassim University, Saudi Arabia

<sup>3</sup>Department of Management Information Systems, College of Business and Economics, Qassim University, Saudi Arabia

**Abstract** In this paper, we define a new metric (distance function) between the dual fuzzy set of hesitation and also between the weighted dual fuzzy set of hesitation. This new measure is made by adding the relative coefficient of variation to the previous measure [53], a comparison is obtained between the previous measure and the present measure. In addition, a comparison is presented between the score function and the distance measure to see which is more efficient and easier to use in decision-making, in the medical diagnosis application. We have proved the triangle inequality of the new normalized Hamming metric, the new Euclidean metric, and the new generalized metric between two dual hesitant fuzzy elements. We have used a Python program to write the code of distance measure calculations. The score function is very important for ordering the dual hesitant fuzzy elements, so it is very important in simplifying the calculations for decision-making.

**Keywords** Distance measure (distance function)- D.F.S.H- W.D.F.S.H.- hesitancy degree- score function - accuracy function- medical diagnosis - decision-making.

**AMS 2010 subject classifications** 46N30-46N60-30F45-03E72

**DOI:** 10.19139/soic-2310-5070-3336

## 1. Introduction

The score function has been widely employed in addressing multi-criteria vague decision-making problems. In its simplest form, the score function is defined as the difference between the membership and non-membership functions [30]. Chen and Tan [2] were among the first to apply this concept in the context of intuitionistic fuzzy sets. Subsequently, De et al. [3] and Kharal [15] proposed three alternative formulations of the S.F. to more accurately capture the predisposition toward positive and negative outcomes. The first defines the score function as the membership degree minus the product of the non-membership and hesitation degrees. The second follows a similar structure but substitutes the product with the arithmetic mean of the non-membership and hesitation degrees. The third defines the score function as the arithmetic mean of the membership and non-membership degrees minus the hesitation degree. Wang [35] used the score function to evaluate the dominated degree, thereby reflecting the relative importance of the criteria in service quality assessment. and when explicitly accounting for hesitation, the S.F. is expressed as the membership degree minus the product of the hesitation and non-membership degrees, which coincides with the first definition introduced by De et al. [3] and Kharal [15].

Although Chen and Tan [2] formalized the score function within fuzzy decision-making, a related notion can be traced back to the cumulative prospect theory (CPT) developed by Tversky and Kahneman [33]. As emphasized

---

\*Correspondence to: Salah Boulaaras (Email: s.boulaaras@qu.edu.sa.). Department of Mathematics, College of Sciences, Qassim University, Saudi Arabia.

by Grabisch et al. [13], CPT provides a decision-making model in which the net predisposition is computed in a straightforward manner as the difference between the positive and negative components, i.e. the membership degree minus the non-membership degree. In recent years, the score function has been successfully applied in a variety of domains, including similarity measures [1], [14], aggregation operators [45], [56], ranking procedures [43], [49], Choquet integrals [29], [47], preference relations [12], [45], programming models [20], [46], multiple-attribute decision making [21], [37], and group decision making [17], [36], [48]. Dual hesitant fuzzy sets provide a powerful means of representing uncertainty in real-world situations by simultaneously employing membership and non-membership degrees. They enable the collection of vague data and their subsequent use in distance-based decision-making tasks [53]. Distance measures, as fundamental constructs in system theory, have numerous applications in diverse fields such as image segmentation, decision making, and artificial intelligence. The distance function, or metric, was extended to Dual hesitant fuzzy sets by Su et al. [27] and has since been applied in a wide range of contexts.

The introduction of fuzzy sets by Zadeh [51] marked a seminal moment in uncertainty modeling, where membership degrees were proposed to capture the complexity of real-world phenomena. This idea had a profound influence and paved the way for subsequent generalizations. In particular, Torra and Narukawa [31,32] generalized the traditional vague set to allow multiple degrees of membership, leading to the development of the hesitant fuzzy set. This framework has since attracted significant scholarly attention. Xia and Xu [38] and Xia et al. [39] studied the mathematical representation of HFS and developed a series of powerful aggregation operators in hesitant environments. Xu and Xia [40] extended entropy and cross-entropy concepts to, deriving several meaningful conclusions. Furthermore, Xu and Xia [41,42] conducted systematic investigations into distance, similarity, and correlation measures in hesitant fuzzy contexts. Farhadinia [10] examined information measures for hesitant fuzzy sets and interval-valued fuzzy sets, while Li et al. [18,19] introduced the concept of hesitancy degree and established distance measures based on it. To broaden the applicability of hesitant fuzzy sets, Zeng et al. [55] proposed a novel distance measure with successful applications in pattern recognition. Collectively, distance and similarity measures for hesitant fuzzy sets have been widely utilized in pattern recognition, approximate reasoning, image segmentation, and medical diagnosis.

The flexibility of vague and hesitant fuzzy systems has also facilitated their use in medical applications. Emanuel et al. [7,8] introduced type-2 fuzzy theory into medical diagnosis, achieving promising results. Similarly, Molla et al. [23] applied Pythagorean fuzzy theory within the Preference Ranking Organization Method for Enrichment Evaluation (PROMETHEE), enhancing diagnostic decision-making.

In practical contexts, dual hesitant fuzzy sets DHFS have proven to be effective tools for managing uncertainty. Singh [26] developed a metric for DHFS and demonstrated its application in evaluating investment alternatives. Su et al. [27] further advanced this line of research by designing distance and similarity measures for DHFS with applications in pattern recognition. However, in certain decision-making scenarios, DHFS may not adequately convey all relevant information to the decision maker. To address this limitation, the weighted dual hesitant fuzzy set (WDHFS) and weighted dual hesitant fuzzy element (WDHFE) were introduced [54]. In this framework, aggregation operators such as the WDHFS averaging operator and the weighted DHFS geometric operator were developed to combine weighted dual hesitant fuzzy information. The concept of a feature vector of a WDHFE was also proposed, consisting of five intrinsic properties: upper bound, lower bound, average level function, standard deviation, and hesitancy degree. Based on this representation, a distance measure was designed to better preserve the original information contained in WDHFE.

Because DHFS incorporates flexible evaluations from experts, it has become a convenient and widely applied framework in decision making and medical diagnosis. Building upon this, Wenyi Zeng et al. [53] introduced several important properties of dual hesitant fuzzy elements (DHFE), including the average function, the variance function, and the degree of hesitancy. They further proposed both standard and weighted metrics for DHFE, with the latter allowing for the adjustment of contributions from different feature indices. Comparative analyses demonstrated the effectiveness of these new distance measures, particularly in capturing medical diagnostic information with greater accuracy. In [11], a novel metric for picture fuzzy sets was introduced, while in [24], a new distance measure for (DFHE) and (WDFHE) was proposed.

The relative coefficient of variation is a powerful tool in medical diagnosis and decision making, it expresses the

degree of variability relative to the mean, making it easier to compare the consistency of tests—even when they use different units or measurement scales. So, we added it to the new distance measure in this paper.[16] In what follows, we will denote the dual fuzzy set of hesitation by D.F.S.H., the weighted dual fuzzy set of hesitation by W.D.F.S.H., the dual fuzzy element of hesitation by D.F.E.H., and the weighted dual fuzzy element of hesitation by W.D.F.E.H. We also denote the score function by S.F.

The structure of the present paper is organized as follows: Part 2. provides the preliminaries of dual hesitant fuzzy sets (DHFS). Introduces key definitions, including the upper and lower bounds, mean function, standard deviation function, coefficient of variation, and hesitancy degree of DHFS.

Part 3. presents the main theoretical contributions and applications. A new distance measure has been formally defined, and several of its properties have been established through rigorous proofs. Furthermore, the effectiveness of the proposed metric is demonstrated in medical diagnosis and decision-making contexts. A comparative analysis between the distance measure and the score function is also provided, highlighting the relative simplicity and practical advantages of the score function. The following chart shows the algorithm used to calculate the distance function in medical diagnosis application:



Figure 1. Chart of the algorithm used in calculating the distance function.

## 2. Preliminaries

**Definition 1.** Let  $U = \{u_1, u_2, \dots, u_n\}$  be the comprehensive set, let  $H$  denote the fuzzy set of hesitation (F.S.H) on  $U$  where

$$H = \{(u, \psi_H(u)) \mid u \in U\} \tag{2.1}$$

and  $\psi_H(u) \subseteq [0, 1]$  is the set of all possible degrees of membership of element  $u \in U$  and let  $\psi_H(u) = \psi(u)$  be the fuzzy element of hesitation (F.E.H.)

For vague sets of hesitation  $\psi(u), \psi_1(u), \psi_2(u)$ , we define the following properties [31,54]:

- (1)  $\psi^- = \min\psi(u), \psi^+(u) = \max\psi(u)$
- (2)  $\bar{\psi}(u) = \cup_{\phi \in \psi} \{1 - \phi\}$
- (3)  $\psi_1(u) \cup \psi_2(u) = \cup_{r_1 \in \psi_1(u), r_2 \in \psi_2(u)} \max\{r_1, r_2\}$
- (4)  $\psi_1(u) \cap \psi_2(u) = \cup_{r_1 \in \psi_1(u), r_2 \in \psi_2(u)} \min\{r_1, r_2\}$
- (5)  $\psi^\lambda(u) = \cup_{r \in \psi} \{r^\lambda\}$
- (6)  $\lambda r(u) = \cup_{r \in \psi} \{1 - (1 - r)^\lambda\}, \lambda > 0$
- (7)  $\psi_1(u) \oplus \psi_2(u) = \cup_{r_1 \in \psi_1(u), r_2 \in \psi_2(u)} \{r_1 + r_2 - r_1 r_2\}$
- (8)  $\psi_1(u) \otimes \psi_2(u) = \cup_{r_1 \in \psi_1(u), r_2 \in \psi_2(u)} \{r_1 r_2\}$

**Definition 2.** Let  $\psi(u)$  be a vague element of hesitation, the score function  $\Upsilon$  of  $\psi(u)$  is defined as follows:

$$\Upsilon(\psi(u)) = \frac{1}{\tau(\psi(u))} \sum_{r \in \psi(u)} r \quad (2.2)$$

where  $\tau(\psi(u))$  is the length of  $\psi(u)$  (the number of elements in  $\psi(u)$ ).

If  $\psi_1(u), \psi_2(u)$  are fuzzy hesitant elements, then

- (i) If  $\Upsilon(\psi_1(u)) \leq \Upsilon(\psi_2(u))$  then  $\psi_1(u) \leq \psi_2(u)$
- (ii) If  $\Upsilon(\psi_1(u)) = \Upsilon(\psi_2(u))$  then  $\psi_1(u) = \psi_2(u)$ .

**Definition 3.** The D.F.S.H. is defined as follows:

$$\Theta = \{(u, \psi(u), \zeta(u)) \mid u \in U\} \quad (2.3)$$

where  $\psi(u) \subseteq [0, 1]$  and  $\zeta(u) \subseteq [0, 1]$  are the sets of all possible membership degree and nonmembership degree of the element  $u \in U$  in the set  $\Theta$ , respectively,  $r \in \psi(u)$  and  $g \in \zeta(u)$  satisfy the following conditions:

- (1)  $r \geq 0, 1 \geq g$
- (2)  $r^+ = \cup_{r \in \psi(u)} \max\{r\}, g^+ = \cup_{g \in \zeta(u)} \max\{g\}$
- (3)  $1 \geq r^+ + g^+ \geq 0$ .

Consider  $\theta = \{\psi(u), \zeta(u)\}$  to be the D.F.E.H.

**Definition 4.** Let  $\theta = \{\psi(u), \zeta(u)\}$  be a dual vague element of hesitation, the score function  $\Upsilon$  of  $\theta$  is defined as follows:

$$\Upsilon(\theta) = \frac{1}{\tau(\psi(u))} \sum_{r \in \psi} r - \frac{1}{\tau(\zeta(u))} \sum_{y \in \zeta} y \quad (2.4)$$

The accuracy function  $k$  of  $\theta$  is defined as follows:

$$k(\theta) = \frac{1}{\tau(\psi(u))} \sum_{r \in \psi} r + \frac{1}{\tau(\zeta(u))} \sum_{y \in \zeta} y \quad (2.5)$$

If  $\theta_1(u), \theta_2(u)$  are D.F.E.H., then if  $\Upsilon(\theta_1(u)) \leq \Upsilon(\theta_2(u))$  then  $\theta_1(u) \leq \theta_2(u)$ ; if  $\Upsilon(\theta_1(u)) = \Upsilon(\theta_2(u))$  then  $\theta_1(u) = \theta_2(u)$ .

**Definition 5.** Let  $\theta^\alpha = \{\psi^\alpha(u), \zeta^\alpha(u)\}$  be a weighted dual hesitant fuzzy element, the score function  $\Upsilon$  of  $\theta$  is defined as follows:

$$\Upsilon(\theta^\alpha) = \sum_{r \in \psi^\alpha} \alpha_r r - \sum_{y \in \zeta^\alpha} \alpha_y y \tag{2.6}$$

The accuracy function  $k$  of  $\theta^\alpha$  is defined as follows:

$$k(\theta^\alpha) = \sum_{r \in \psi^\alpha} \alpha_r r + \sum_{y \in \zeta^\alpha} \alpha_y y \tag{2.7}$$

If  $\theta_1^\alpha(u), \theta_2^\alpha(u)$  are W.F.E.H. that are hesitant, then if  $\Upsilon(\theta_1^\alpha(u)) \leq \Upsilon(\theta_2^\alpha(u))$  then  $\theta_1^\alpha(u) \leq \theta_2^\alpha(u)$ ; if  $\Upsilon(\theta_1^\alpha(u)) = \Upsilon(\theta_2^\alpha(u))$  then  $\theta_1^\alpha(u) = \theta_2^\alpha(u)$  if  $k(\theta_1^\alpha) = k(\theta_2^\alpha)$  and  $\theta_1^\alpha(u) \leq \theta_2^\alpha(u)$  if  $k(\theta_1^\alpha) \leq k(\theta_2^\alpha)$

**Definition 6.** Let  $O = (o_{ij})_{(m \times n)}$  be the collective intuitionistic fuzzy decision matrix; then we call  $S = (s_{ij})_{(m \times n)}$  the score matrix of  $O = (o_{ij})_{(m \times n)}$  where:

$$s_i = s(o_{ij}) = \psi(ij) - \zeta(ij), i = 1, 2, \dots, m; j = 1, 2, \dots, n \tag{2.8}$$

and  $s(o_{ij})$  is called the score of  $o_{ij}$

The score value of  $s_{ij}$  is defined as follows:

$$s_j(w) = \sum_{i=1}^m w_i s_{ij} \tag{2.9}$$

**Definition 7.(Distance measure)** The function  $\delta(DHFS, DHFS)$  is called a distance measure or metric between E, F, if the following conditions hold:

- (1)  $1 \geq \delta(L, M) \geq 0$ .
- (2)  $\delta(L, M) = 0$  if and only if  $E = F$ .
- (3)  $\delta(L, M) = \delta(M, L)$ .
- (4)  $\delta(L, M) \leq \delta(L, N) + \delta(N, M)$ .

where  $L = \{(u, \psi_L(u), \zeta_L(u)); u \in U\}$ ,  $M = \{(u, \psi_M(u), \zeta_M(u)); u \in U\}$ ,  $N = \{(u, \psi_N(u), \zeta_N(u)); u \in U\}$  are D.F.S.H. and  $\delta(L, M) : DHFS \times DHFS \rightarrow [0, 1]$ .

**Definition 8.** Let  $\theta = \{\psi, \zeta\}$  then  $\theta^+ = \{\theta_\psi^+, \theta_\zeta^+\}$   
 $= \{\max\{r\}_{r \in \psi}, \min\{u\}_{u \in \zeta}\}$ ,  $\theta^- = \{\theta_\psi^-, \theta_\zeta^-\} = \{\min\{r\}_{r \in \psi}, \max\{r\}_{u \in \zeta}\}$  are called the upper and lower bound of  $\theta$ , respectively.

**Definition 9.** If  $\theta = \{\psi, \zeta\}$  then

$$\phi(\theta) = \{\phi_\psi, \phi_\zeta\} = \left\{ \frac{1}{\tau(\psi)} \sum_{r \in \psi} r, \frac{1}{\tau(\zeta)} \sum_{u \in \zeta} u \right\} \tag{2.10}$$

and

$$b(\theta) = \{b_\psi, b_\zeta\} = \left\{ \sqrt{\frac{1}{\tau(\psi)} \sum_{r \in \psi} (r - \phi_\psi)^2}, \sqrt{\frac{1}{\tau(\zeta)} \sum_{u \in \zeta} (u - \phi_\zeta)^2} \right\} \quad (2.11)$$

and

$$c(\theta) = \{c_\psi, c_k\} = \left\{ \frac{\sqrt{\frac{1}{\tau(\psi)} \sum_{r \in \psi} (r - \phi_\psi)^2}}{\frac{1}{\tau(\psi)} \sum_{r \in \psi} r}, \frac{\sqrt{\frac{1}{\tau(\zeta)} \sum_{u \in \zeta} (u - \phi_\zeta)^2}}{\frac{1}{\tau(\zeta)} \sum_{u \in \zeta} u} \right\} \quad (2.12)$$

denote the mean function, the standard deviation, and the relative coefficient of variation of the F.E.H. respectively.

$\tau(\psi), \tau(\zeta)$  are the lengths of  $\psi, \zeta$  (the number of elements in each)

$$\eta(\theta) = \{\eta_\psi, \eta_\zeta\} = \left\{ 1 - \frac{1}{\tau(\psi)}, 1 - \frac{1}{\tau(k)} \right\} \quad (2.13)$$

denotes the degree of hesitancy of  $\theta$ .

### 3. The Fundamental Results

#### 3.1. The New Distance Measure

**Definition 10.** If  $\theta_1$  and  $\theta_2$  are two D.F.E, then

$$\begin{aligned} \delta_1(\theta_1, \theta_2) = \frac{1}{12} & \left( |\theta_{1\psi}^+ - \theta_{2\psi}^+| + |\theta_{1\psi}^- - \theta_{2\psi}^-| + |\theta_{1\zeta}^+ - \theta_{2\zeta}^+| + |\theta_{1\zeta}^- - \theta_{2\zeta}^-| \right. \\ & + |\phi_{1\psi} - \phi_{2\psi}| + |\phi_{1\zeta} - \phi_{2\zeta}| + |b_{1\psi} - b_{2\psi}| + |b_{1\zeta} - b_{2\zeta}| \\ & \left. + |\eta_{1\psi} - \eta_{2\psi}| + |\eta_{1\zeta} - \eta_{2\zeta}| + |c_{1\psi} - c_{2\psi}| + |c_{1\zeta} - c_{2\zeta}| \right) \end{aligned} \quad (3.1)$$

$$\begin{aligned} \delta_2(\theta_1, \theta_2) = & \left[ \frac{1}{12} \left( (\theta_{1\psi}^+ - \theta_{2\psi}^+)^2 + (\theta_{1\psi}^- - \theta_{2\psi}^-)^2 + (\theta_{1\zeta}^+ - \theta_{2\zeta}^+)^2 + (\theta_{1\zeta}^- - \theta_{2\zeta}^-)^2 \right. \right. \\ & + (\phi_{1\psi} - \phi_{2\psi})^2 + (\phi_{1\zeta} - \phi_{2\zeta})^2 + (b_{1\psi} - b_{2\psi})^2 + (b_{1\zeta} - b_{2\zeta})^2 \\ & \left. \left. + (\eta_{1\psi} - \eta_{2\psi})^2 + (\eta_{1\zeta} - \eta_{2\zeta})^2 + (c_{1\psi} - c_{2\psi})^2 + (c_{1\zeta} - c_{2\zeta})^2 \right) \right]^{1/2} \end{aligned} \quad (3.2)$$

$$\begin{aligned} \delta_3(\theta_1, \theta_2) = & \left[ \frac{1}{12} \left( |\theta_{1\psi}^+ - \theta_{2\psi}^+|^\lambda + |\theta_{1\psi}^- - \theta_{2\psi}^-|^\lambda + |\theta_{1\zeta}^+ - \theta_{2\zeta}^+|^\lambda + |\theta_{1\zeta}^- - \theta_{2\zeta}^-|^\lambda \right. \right. \\ & + |\phi_{1\psi} - \phi_{2\psi}|^\lambda + |\phi_{1\zeta} - \phi_{2\zeta}|^\lambda + |b_{1\psi} - b_{2\psi}|^\lambda + |b_{1\zeta} - b_{2\zeta}|^\lambda \\ & \left. \left. + |\eta_{1\psi} - \eta_{2\psi}|^\lambda + |\eta_{1\zeta} - \eta_{2\zeta}|^\lambda + |c_{1\psi} - c_{2\psi}|^\lambda + |c_{1\zeta} - c_{2\zeta}|^\lambda \right) \right]^{\frac{1}{\lambda}} \end{aligned} \quad (3.3)$$

where  $\lambda > 0$ ,  $\delta_1(\theta_1, \theta_2)$  is the normalized Hamming metric,  $\delta_2(\theta_1, \theta_2)$  is the normalized Euclidean metric, and  $\delta_3(\theta_1, \theta_2)$  is the normalized generalized metric between  $\theta_1$  and  $\theta_2$ .

**Theorem 1.** Let  $\theta_1, \theta_2$  be two dual hesitant fuzzy elements then:

- (i) Normalized Hamming metric  $\delta_1(\theta_1, \theta_2)$
  - (ii) Normalized Euclidean metric  $\delta_2(\theta_1, \theta_2)$
  - (iii) Normalized generalized metric  $\delta_3(\theta_1, \theta_2)$
- satisfy the conditions of distance measurement.

**Proof.** (i) The normalized Hamming metric  $\delta_1(\theta_1, \theta_2)$  satisfies the following conditions:

- (1)  $1 \geq \delta_1(\theta_1, \theta_2) \geq 0$ .
- (2)  $\delta_1(\theta_1, \theta_2) = 0$  if and only if  $\theta_1 = \theta_2$ .
- (3)  $\delta_1(\theta_1, \theta_2) = \delta_1(\theta_2, \theta_1)$ .
- (4)

$$\begin{aligned} \delta_1(\theta_1, \theta_3) &= \frac{1}{12} (|\theta_{1\psi}^+ - \theta_{3\psi}^+| + |\theta_{1\psi}^- - \theta_{3\psi}^-| + |\theta_{1\zeta}^+ - \theta_{3\zeta}^+| + |\theta_{1\zeta}^- - \theta_{3\zeta}^-| + |\phi_{1\psi} - \phi_{3\psi}| \\ &\quad + |\phi_{1\zeta} - \phi_{3\zeta}| + |b_{1\psi} - b_{3\psi}| + |b_{1\zeta} - b_{3\zeta}| + |\eta_{1\psi} - \eta_{3\psi}| + |\eta_{1\zeta} - \eta_{3\zeta}| \\ &\quad + |c_{1\psi} - c_{2\psi}| + |c_{1\zeta} - c_{2\zeta}|) \\ &\leq \frac{1}{12} (|\theta_{1\psi}^+ - \theta_{2\psi}^+| + |\theta_{1\psi}^- - \theta_{2\psi}^-| + |\theta_{1\zeta}^+ - \theta_{2\zeta}^+| + |\theta_{1\zeta}^- - \theta_{2\zeta}^-| + |\phi_{1\psi} - \phi_{2\psi}| \\ &\quad + |\phi_{1\zeta} - \phi_{2\zeta}| + |b_{1\psi} - b_{2\psi}| + |b_{1\zeta} - b_{2\zeta}| + |\eta_{1\psi} - \eta_{2\psi}| + |\eta_{1\zeta} - \eta_{2\zeta}| \\ &\quad + |c_{1\psi} - c_{2\psi}| + |c_{1\zeta} - c_{2\zeta}|) \\ &\quad + \frac{1}{12} (|\theta_{2\psi}^+ - \theta_{3\psi}^+| + |\theta_{2\psi}^- - \theta_{3\psi}^-| + |\theta_{2\zeta}^+ - \theta_{3\zeta}^+| + |\theta_{2\zeta}^- - \theta_{3\zeta}^-| + |\phi_{2\psi} - \phi_{3\psi}| \\ &\quad + |\phi_{2\zeta} - \phi_{3\zeta}| + |b_{2\psi} - b_{3\psi}| + |b_{2\zeta} - b_{3\zeta}| + |\eta_{2\psi} - \eta_{3\psi}| + |\eta_{2\zeta} - \eta_{3\zeta}| \\ &\quad + |c_{1\psi} - c_{2\psi}| + |c_{1\zeta} - c_{2\zeta}|) \\ &= \delta_1(\theta_1, \theta_2) + \rho_1(\theta_2, \theta_3) \end{aligned}$$

(ii) The normalized Euclidean distance measure  $\rho_2(\theta_1, \theta_2)$  also satisfies the axioms of the distance measure:

- (1)  $1 \geq \delta_2(\theta_1, \theta_2) \geq 0$ .
- (2)  $\delta_2(\theta_1, \theta_2) = 0$  if and only if  $\theta_1 = \theta_2$ .
- (3)  $\delta_2(\theta_1, \theta_2) = \delta_2(\theta_2, \theta_1)$ .
- (4)

$$\begin{aligned} \delta_2(\theta_1, \theta_3) &= \left( \frac{1}{12} (d_{1\psi}^+ - \theta_{3\psi}^+)^2 + (\theta_{1\psi}^- - \theta_{3\psi}^-)^2 + (\theta_{1\zeta}^+ - \theta_{3\zeta}^+)^2 + (\theta_{1\zeta}^- - \theta_{3\zeta}^-)^2 + (\phi_{1\psi} - \phi_{3\psi})^2 \right. \\ &\quad + (g_{1\zeta} - g_{3\zeta})^2 + (b_{1\psi} - b_{3\psi})^2 + (b_{1\zeta} - b_{3\zeta})^2 + (\eta_{1\psi} - \eta_{3\psi})^2 + (\eta_{1\zeta} - \eta_{3\zeta})^2 \\ &\quad \left. (c_{1\psi} - c_{3\psi})^2 + (c_{1\zeta} - c_{3\zeta})^2 \right)^{1/2} \\ &\leq \left( \frac{1}{12} (\theta_{1\psi}^+ - \theta_{2\psi}^+)^2 + (\theta_{1\psi}^- - \theta_{2\psi}^-)^2 + (\theta_{1\zeta}^+ - \theta_{2\zeta}^+)^2 + (\theta_{1\zeta}^- - \theta_{2\zeta}^-)^2 + (\phi_{1\psi} - \phi_{2\psi})^2 \right. \\ &\quad + (\phi_{1\zeta} - \phi_{2\zeta})^2 + (b_{1\psi} - b_{2\psi})^2 + (b_{1\zeta} - b_{2\zeta})^2 + (\eta_{1\psi} - \eta_{2\psi})^2 + (\eta_{1\zeta} - \eta_{2\zeta})^2 \\ &\quad \left. (c_{1\psi} - c_{2\psi})^2 + (c_{1\zeta} - c_{2\zeta})^2 \right)^{1/2} \end{aligned}$$

$$\begin{aligned}
& + \left( \frac{1}{12} (\theta_{2\psi}^+ - \theta_{3\psi}^+)^2 + (\theta_{2\psi}^- - \theta_{3\psi}^-)^2 + (\theta_{2\zeta}^+ - \theta_{3\zeta}^+)^2 + (\theta_{2\zeta}^- - \theta_{3\zeta}^-)^2 + (\phi_{2\psi} - \phi_{3\psi})^2 \right. \\
& \quad + (\phi_{2\zeta} - \phi_{3\zeta})^2 + (b_{2\psi} - b_{3\psi})^2 + (b_{2\zeta} - b_{3\zeta})^2 + (\eta_{2\psi} - \eta_{3\psi})^2 + (\eta_{2\zeta} - \eta_{3\zeta})^2 \\
& \quad \left. (c_{2\psi} - c_{3\psi})^2 + (c_{2\zeta} - c_{3\zeta})^2 \right)^{1/2} \\
& = \rho_2(\theta_1, \theta_2) + \rho_2(\theta_2, \theta_3)
\end{aligned}$$

(iii) The metric axioms hold for the normalized generalized metric  $\rho_3(\theta_1, \theta_2)$  since:

- (1)  $1 \geq \delta_3(\theta_1, \theta_2) \geq 0$ .
- (2)  $\delta_3(\theta_1, \theta_2) = 0$  if and only if  $\theta_1 = \theta_2$ .
- (3)  $\delta_3(\theta_1, \theta_2) = \delta_1(\theta_2, \theta_1)$ .
- (4)

$$\begin{aligned}
\delta_3(\theta_1, \theta_3) &= \left( \frac{1}{12} (|\theta_{1\psi}^+ - \theta_{3\psi}^+|^\lambda + |\theta_{1\psi}^- - \theta_{3\psi}^-|^\lambda + |\theta_{1\zeta}^+ - \theta_{3\zeta}^+|^\lambda + |\theta_{1\zeta}^- - \theta_{3\zeta}^-|^\lambda + |\phi_{1\psi} - \phi_{3\psi}|^\lambda \right. \\
& \quad + |\phi_{1\zeta} - \phi_{3\zeta}|^\lambda + |b_{1\psi} - b_{3\psi}|^\lambda + |b_{1\zeta} - b_{3\zeta}|^\lambda + |\eta_{1\psi} - \eta_{3\psi}|^\lambda + |\eta_{1\zeta} - \eta_{3\zeta}|^\lambda \\
& \quad \left. + |c_{1\psi} - c_{3\psi}|^\lambda + |c_{1\zeta} - c_{3\zeta}|^\lambda \right)^{1/\lambda} \\
&\leq \left( \frac{1}{12} (|\theta_{1\psi}^+ - \theta_{2\psi}^+|^\lambda + |\theta_{1\psi}^- - \theta_{2\psi}^-|^\lambda + |\theta_{1\zeta}^+ - \theta_{2\zeta}^+|^\lambda + |\theta_{1\zeta}^- - \theta_{2\zeta}^-|^\lambda + |\phi_{1\psi} - \phi_{2\psi}|^\lambda \right. \\
& \quad + |\phi_{1\zeta} - \phi_{2\zeta}|^\lambda + |b_{1\psi} - b_{2\psi}|^\lambda + |b_{1\zeta} - b_{2\zeta}|^\lambda + |\eta_{1\psi} - \eta_{2\psi}|^\lambda + |\eta_{1\zeta} - \eta_{2\zeta}|^\lambda \\
& \quad \left. + |c_{1\psi} - c_{2\psi}|^\lambda + |c_{1\zeta} - c_{2\zeta}|^\lambda \right)^{1/\lambda} \\
&+ \left( \frac{1}{12} (|\theta_{2\psi}^+ - \theta_{3\psi}^+|^\lambda + |\theta_{2\psi}^- - \theta_{3\psi}^-|^\lambda + |\theta_{2\zeta}^+ - \theta_{3\zeta}^+|^\lambda + |\theta_{2\zeta}^- - \theta_{3\zeta}^-|^\lambda + |\phi_{2\psi} - \phi_{3\psi}|^\lambda \right. \\
& \quad + |\phi_{2\zeta} - \phi_{3\zeta}|^\lambda + |b_{2\psi} - b_{3\psi}|^\lambda + |b_{2\zeta} - b_{3\zeta}|^\lambda + |\eta_{2\psi} - \eta_{3\psi}|^\lambda + |\eta_{2\zeta} - \eta_{3\zeta}|^\lambda \\
& \quad \left. + |c_{2\psi} - c_{3\psi}|^\lambda + |c_{2\zeta} - c_{3\zeta}|^\lambda \right)^{1/\lambda} \\
&= \delta_3(\theta_1, \theta_2) + \delta_3(\theta_2, \theta_3)
\end{aligned}$$

□

The weighted normalized Hamming metric, the weighted normalized Euclidean metric, and the weighted normalized generalized metric between  $\theta_1$  and  $\theta_2$  are defined as follows:

$$\begin{aligned}
\delta_\alpha^1(\theta_1, \theta_2) &= \alpha_1 |\theta_{1\psi}^+ - \theta_{2\psi}^+| + \alpha_2 |\theta_{1\psi}^- - \theta_{2\psi}^-| + \alpha_3 |\theta_{1\zeta}^+ - \theta_{2\zeta}^+| + \alpha_4 |\theta_{1\zeta}^- - \theta_{2\zeta}^-| + \alpha_5 |\phi_{1\psi} - \phi_{2\psi}| \\
& \quad + \alpha_6 |\phi_{1\zeta} - \phi_{2\zeta}| + \alpha_7 |b_{1\psi} - b_{2\psi}| + \alpha_8 |b_{1\zeta} - b_{2\zeta}| + \alpha_9 |\eta_{1\psi} - \eta_{2\psi}| + \alpha_{10} |\eta_{1\zeta} - \eta_{2\zeta}| \\
& \quad + \alpha_{11} |c_{1\psi} - c_{2\psi}| + \alpha_{12} |c_{1\zeta} - c_{2\zeta}|
\end{aligned} \tag{3.4}$$

$$\begin{aligned}
\delta_\alpha^2(\theta_1, \theta_2) &= (\alpha_1 (\theta_{1\psi}^+ - \theta_{2\psi}^+)^2 + \alpha_2 (\theta_{1\psi}^- - \theta_{2\psi}^-)^2 + \alpha_3 (\theta_{1\zeta}^+ - \theta_{2\zeta}^+)^2 + \alpha_4 (\theta_{1\zeta}^- - \theta_{2\zeta}^-)^2 \\
& \quad + \alpha_5 (\phi_{1\psi} - \phi_{2\psi})^2 + \alpha_6 (\phi_{1\zeta} - \phi_{2\zeta})^2 + \alpha_7 (b_{1\psi} - b_{2\psi})^2 \\
& \quad + \alpha_8 (b_{1\zeta} - b_{2\zeta})^2 + \alpha_9 (\eta_{1\psi} - \eta_{2\psi})^2 + \alpha_{10} (\eta_{1\zeta} - \eta_{2\zeta})^2)^{1/2} \\
& \quad + \alpha_{11} (c_{1\psi} - c_{2\psi})^2 + \alpha_{12} (c_{1\zeta} - c_{2\zeta})^2)^{1/2}
\end{aligned} \tag{3.5}$$

$$\begin{aligned} \delta_{\alpha}^3(\theta_1, \theta_2) = & ((\alpha_1|\theta_{1\psi}^+ - \theta_{2\psi}^+|^{\lambda} + \alpha_2|\theta_{1\psi}^- - \theta_{2\psi}^-|^{\lambda} + \alpha_3|\theta_{1\zeta}^+ - \theta_{2\zeta}^+|^{\lambda} + \alpha_4|d_{1\zeta}^- - d_{2\zeta}^-|^{\lambda} \\ & + \alpha_5|\phi_{1\psi} - \phi_{2\psi}|^{\lambda} + \alpha_6|\phi_{1\zeta} - \phi_{2\zeta}|^{\lambda} + \alpha_7|b_{1\psi} - b_{2\psi}|^{\lambda} \\ & + \alpha_8|b_{1\zeta} - b_{2\zeta}|^{\lambda} + \alpha_9|\eta_{1\psi} - \eta_{2\psi}|^{\lambda} + \alpha_{10}|\eta_{1\zeta} - \eta_{2\zeta}|^{\lambda}))^{1/\lambda} \\ & + \alpha_{11}|c_{1\psi} - c_{2\psi}|^{\lambda} + \alpha_{12}|c_{1\zeta} - c_{2\zeta}|^{\lambda})^{1/\lambda} \end{aligned} \quad (3.6)$$

where  $\alpha_i$  are the weights.

**Theorem 2.** Let  $\theta_1, \theta_2$  be two W.D.F.E.H then the weighted normalized Hamming metric, the weighted normalized Euclidean metric and the weighted normalized generalized metric satisfy the triangle inequality.

**Proof.:** The proof of this theorem is similar to theorem 1.  $\square$

**Definition 11.** Let  $U = \{u_1, u_2, \dots, u_n\}$  and  $\Theta_1$  and  $\Theta_2$  dual hesitant fuzzy sets on  $U$  then we can define the distance function between  $\Theta_1$  and  $\Theta_2$  as follow

$$\begin{aligned} \delta_1(\Theta_1, \Theta_2) = & \frac{1}{12n} \sum_{i=1}^n (|\theta_{1\psi}^+ - \theta_{2\psi}^+| + |\theta_{1\psi}^- - \theta_{2\psi}^-| + |\theta_{1\zeta}^+ - \theta_{2\zeta}^+| + |\theta_{1\zeta}^- - \theta_{2\zeta}^-| + |\phi_{1\psi} - \phi_{2\psi}| \\ & + |\phi_{1\zeta} - \phi_{2\zeta}| + |b_{1\psi} - b_{2\psi}| + |b_{1\zeta} - b_{2\zeta}| + |\eta_{1\psi} - \eta_{2\psi}| + |\eta_{1\zeta} - \eta_{2\zeta}| \\ & + |c_{1\psi} - c_{2\psi}| + |c_{1\zeta} - c_{2\zeta}|) \end{aligned} \quad (3.7)$$

$$\begin{aligned} \delta_2(\Theta_1, \Theta_2) = & \left( \frac{1}{12n} \sum_{i=1}^n (\theta_{1\psi}^+ - \theta_{2\psi}^+)^2 + (\theta_{1\psi}^- - \theta_{2\psi}^-)^2 + (\theta_{1\zeta}^+ - \theta_{2\zeta}^+)^2 + (\theta_{1\zeta}^- - \theta_{2\zeta}^-)^2 + (\phi_{1\psi} - \phi_{2\psi})^2 \right. \\ & + (\phi_{1\zeta} - \phi_{2\zeta})^2 + (b_{1\psi} - b_{2\psi})^2 + (b_{1\zeta} - b_{2\zeta})^2 + (\eta_{1\psi} - \eta_{2\psi})^2 + (\eta_{1\zeta} - \eta_{2\zeta})^2 \\ & \left. + (c_{1\psi} - c_{2\psi})^2 + (c_{1\zeta} - c_{2\zeta})^2 \right)^{1/2} \end{aligned} \quad (3.8)$$

$$\begin{aligned} \delta_3(\Theta_1, \Theta_2) = & \left( \frac{1}{12n} \sum_{i=1}^n (|\theta_{1\psi}^+ - \theta_{2\psi}^+|^{\lambda} + |\theta_{1\psi}^- - \theta_{2\psi}^-|^{\lambda} + |\theta_{1\zeta}^+ - \theta_{2\zeta}^+|^{\lambda} + |\theta_{1\zeta}^- - \theta_{2\zeta}^-|^{\lambda} + |\phi_{1\psi} - \phi_{2\psi}|^{\lambda} \right. \\ & + |\phi_{1\zeta} - \phi_{2\zeta}|^{\lambda} + |b_{1\psi} - b_{2\psi}|^{\lambda} + |b_{1\zeta} - b_{2\zeta}|^{\lambda} + |\eta_{1\psi} - \eta_{2\psi}|^{\lambda} + |\eta_{1\zeta} - \eta_{2\zeta}|^{\lambda} \\ & \left. + |c_{1\psi} - c_{2\psi}|^{\lambda} + |c_{1\zeta} - c_{2\zeta}|^{\lambda}) \right)^{1/\lambda} \end{aligned} \quad (3.9)$$

where  $\lambda > 0$ ,  $\delta_1(\Theta_1, \Theta_2)$  is the normalized Hamming metric,  $\delta_2(\Theta_1, \Theta_2)$  is the normalized Euclidean metric, and  $\delta_3(\Theta_1, \Theta_2)$  is the normalized generalized metric between D.F.S.H.  $\Theta_1$  and  $\Theta_2$  depending on the axioms of D.F.E.H.

**Definition 12.** Let  $\Theta_1$  and  $\Theta_2$  be D.F.S.H. on  $U = \{u_1, u_2, \dots, u_n\}$ , the weighted normalized Hamming metric, the weighted normalized Euclidean metric and the weighted normalized generalized

metric between  $\Theta_1$  and  $\Theta_2$  are defined, respectively, as follows:

$$\begin{aligned} \delta_{\alpha}^1(\Theta_1, \Theta_2) = & \frac{1}{n} \sum_{i=1}^n (\alpha_1 |\theta_{1\psi}^+ - \theta_{2\psi}^+| + \alpha_2 |\theta_{1\psi}^- - \theta_{2\psi}^-| + \alpha_3 |\theta_{1\zeta}^+ - \theta_{2\zeta}^+| + \alpha_4 |\theta_{1\zeta}^- - \theta_{2\zeta}^-| + \alpha_5 |\phi_{1\psi} - \phi_{2\psi}| \\ & + \alpha_6 |\phi_{1\zeta} - \phi_{2\zeta}| + \alpha_7 |b_{1\psi} - b_{2\psi}| + \alpha_8 |b_{1\zeta} - b_{2\zeta}| + \alpha_9 |\eta_{1\psi} - \eta_{2\psi}| + \alpha_{10} |\eta_{1\zeta} - \eta_{2\zeta}| \\ & + \alpha_{11} |c_{1\psi} - c_{2\psi}| + \alpha_{12} |c_{1\zeta} - c_{2\zeta}|) \end{aligned} \quad (3.10)$$

$$\begin{aligned} \delta_{\alpha}^2(\Theta_1, \Theta_2) = & \left( \frac{1}{n} \sum_{i=1}^n (\alpha_1 (\theta_{1\psi}^+ - \theta_{2\psi}^+)^2 + \alpha_2 (\theta_{1\psi}^- - \theta_{2\psi}^-)^2 + \alpha_3 (\theta_{1\zeta}^+ - \theta_{2\zeta}^+)^2 + \alpha_4 (\theta_{1\zeta}^- - \theta_{2\zeta}^-)^2 \right. \\ & + \alpha_5 (\phi_{1\psi} - \phi_{2\psi})^2 + \alpha_6 (\phi_{1\zeta} - \phi_{2\zeta})^2 + \alpha_7 (b_{1\psi} - b_{2\psi})^2 \\ & + \alpha_8 (b_{1\zeta} - b_{2\zeta})^2 + \alpha_9 (\eta_{1\psi} - \eta_{2\psi})^2 + \alpha_{10} (\eta_{1\zeta} - \eta_{2\zeta})^2 \\ & \left. + \alpha_{11} (c_{1\psi} - c_{2\psi})^2 + \alpha_{12} (c_{1\zeta} - c_{2\zeta})^2 \right)^{1/2} \end{aligned} \quad (3.11)$$

$$\begin{aligned} \delta_{\alpha}^3(\Theta_1, \Theta_2) = & \left( \frac{1}{12n} \sum_{i=1}^n (\alpha_1 |\theta_{1\psi}^+ - \theta_{2\psi}^+|^{\lambda} + \alpha_2 |\theta_{1\psi}^- - \theta_{2\psi}^-|^{\lambda} + \alpha_3 |\theta_{1\zeta}^+ - \theta_{2\zeta}^+|^{\lambda} + \alpha_4 |\theta_{1\zeta}^- - \theta_{2\zeta}^-|^{\lambda} \right. \\ & + \alpha_5 |\phi_{1\psi} - \phi_{2\psi}|^{\lambda} + \alpha_6 |\phi_{1\zeta} - \phi_{2\zeta}|^{\lambda} + \alpha_7 |b_{1\psi} - b_{2\psi}|^{\lambda} \\ & + \alpha_8 |b_{1\zeta} - b_{2\zeta}|^{\lambda} + \alpha_9 |\eta_{1\psi} - \eta_{2\psi}|^{\lambda} + \alpha_{10} |\eta_{1\zeta} - \eta_{2\zeta}|^{\lambda} \\ & \left. + \alpha_{11} |c_{1\psi} - c_{2\psi}|^{\lambda} + \alpha_{12} |c_{1\zeta} - c_{2\zeta}|^{\lambda} \right)^{1/\lambda} \end{aligned} \quad (3.12)$$

These definitions are based on the properties of D.F.S.H. where  $0 \leq \alpha_i \leq 1$  and  $\sum_{i=1}^{12} \alpha_i = 1$ .

### 3.2. Motivation Of The New Distance Measure

The previous distance measure in[53] is defined by the following equation:

$$\begin{aligned} \delta_1(\Theta_1, \Theta_2) = & \frac{1}{10n} \sum_{i=1}^n (|\theta_{1\psi}^+ - \theta_{2\psi}^+| + |\theta_{1\psi}^- - \theta_{2\psi}^-| + |\theta_{1\zeta}^+ - \theta_{2\zeta}^+| + |\theta_{1\zeta}^- - \theta_{2\zeta}^-| + |\phi_{1\psi} - \phi_{2\psi}| \\ & + |\phi_{1\zeta} - \phi_{2\zeta}| + |b_{1\psi} - b_{2\psi}| + |b_{1\zeta} - b_{2\zeta}| + |\eta_{1\psi} - \eta_{2\psi}| + |\eta_{1\zeta} - \eta_{2\zeta}|) \end{aligned} \quad (3.13)$$

Existing distance measures for dual hesitant fuzzy sets (3-14) incorporate the mean, variance, and hesitancy degree. However, these quantities are treated independently and fail to capture the interaction between central tendency and dispersion. In particular, variance is scale-dependent and does not adequately reflect relative uncertainty. For example, consider two hesitant fuzzy elements  $q_1 = 0.1, 0.9$  and  $q_2 = 0.49, 0.51$ . Both have the same mean, but  $q_1$  exhibits significantly higher dispersion than  $q_2$ . Classical distance measures assign them nearly equal distances, failing to distinguish between highly inconsistent and highly consistent information. Moreover, the hesitancy degree only reflects the number of elements and ignores how widely the values are distributed. To overcome these limitations, we introduce the relative coefficient of variation, which normalizes the standard deviation by the mean and captures relative dispersion. This allows the proposed distance measure to distinguish between

alternatives with similar averages but different levels of uncertainty. Consequently, the proposed distance provides a more robust and conceptually complete representation of dual hesitant fuzzy information, while remaining consistent with classical measures when dispersion is small.

**3.3. The Applications And The Comparative Work**

**Example 1** [50, 53] In this application, we discuss how to make a decision using the score function and show how easy it is to do this. A car factory needs to decide the best global supplier for one of its most important parts used in collection operation. The characteristics considered here in the selection of five global suppliers  $M_i, i = \{1, 2, 3, 4, 5\}$  are: (1)  $p_1$  is the cost of the product; (2)  $p_2$  is the quality of the product; (3)  $p_3$  is the service performance of the supplier; (4)  $p_4$  is the supplier’s profile and (5)  $p_5$  is a risk factor.  $z = (0.2, 0.15, 0.2, 0.3, 0.15)^T$  is the weight vector of  $S_j$ .  $\omega = (0.4, 0.2, 0.1, 0.3)^T$  is the weight vector of the experts group  $S_l(l = 1, 2, 3, 4)$  which consists of four specialists from each strategic area. The five global suppliers  $M_1, M_2, M_3, M_4, M_5$  will be evaluated using fuzzy intuition numbers by the specialists.

four intuitionistic fuzzy decision matrices are made by specialists, which denotes the satisfaction and dissatisfaction of specialists or experts  $S_l(l = 1, 2, 3, 4)$  for the possible values of the supplier  $M_i$  with respect to the properties  $p_j$ .

The following tables show the W.D.F.S.H., made by experts:

Table 1. Intuitionistic fuzzy decision matrix  $S^1$

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$p_1$	(0.8, 0.1)	(0.5, 0.1)	(0.5, 0.5)	(0.7, 0.1)	(0.8, 0.1)
$p_2$	(0.6, 0.2)	(0.8, 0.2)	(0.6, 0.3)	(0.6, 0.1)	(0.4, 0.5)
$p_3$	(0.7, 0.3)	(0.3, 0.4)	(0.3, 0.2)	(0.9, 0.1)	(0.7, 0.2)
$p_4$	(0.5, 0.2)	(0.7, 0.2)	(0.7, 0.3)	(0.4, 0.3)	(0.5, 0.2)
$p_5$	(0.4, 0.5)	(0.6, 0.2)	(0.8, 0.1)	(0.3, 0.4)	(0.3, 0.4)

Table 2. Intuitionistic fuzzy decision matrix  $S^2$

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$p_1$	(0.7, 0.3)	(0.7, 0.2)	(0.4, 0.2)	(0.7, 0.3)	(0.5, 0.1)
$p_2$	(0.6, 0.1)	(0.6, 0.2)	(0.6, 0.3)	(0.7, 0.2)	(0.6, 0.2)
$p_3$	(0.8, 0.2)	(0.4, 0.4)	(0.7, 0.2)	(0.4, 0.2)	(0.5, 0.4)
$p_4$	(0.7, 0.1)	(0.6, 0.2)	(0.6, 0.1)	(0.8, 0.1)	(0.6, 0.3)
$p_5$	(0.5, 0.3)	(0.7, 0.3)	(0.5, 0.3)	(0.5, 0.4)	(0.7, 0.3)

Table 3. Intuitionistic fuzzy decision matrix  $S^3$

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$p_1$	(0.7, 0.3)	(0.8, 0.2)	(0.8, 0.2)	(0.6, 0.1)	(0.7, 0.1)
$p_2$	(0.8, 0.1)	(0.5, 0.3)	(0.6, 0.1)	(0.5, 0.4)	(0.6, 0.2)
$p_3$	(0.6, 0.4)	(0.5, 0.2)	(0.7, 0.3)	(0.9, 0.1)	(0.8, 0.1)
$p_4$	(0.5, 0.2)	(0.6, 0.3)	(0.4, 0.2)	(0.5, 0.2)	(0.7, 0.3)
$p_5$	(0.6, 0.3)	(0.6, 0.4)	(0.8, 0.1)	(0.6, 0.3)	(0.5, 0.2)

Table 4. Intuitionistic fuzzy decision matrix  $S^4$

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$p_1$	(0.9, 0.1)	(0.7, 0.3)	(0.8, 0.1)	(0.7, 0.2)	(0.8, 0.1)
$p_2$	(0.8, 0.1)	(0.7, 0.1)	(0.6, 0.2)	(0.6, 0.4)	(0.6, 0.3)
$p_3$	(0.5, 0.5)	(0.3, 0.1)	(0.6, 0.3)	(0.7, 0.2)	(0.8, 0.2)
$p_4$	(0.4, 0.6)	(0.8, 0.2)	(0.4, 0.1)	(0.5, 0.1)	(0.7, 0.2)
$p_5$	(0.3, 0.4)	(0.4, 0.2)	(0.5, 0.2)	(0.8, 0.2)	(0.6, 0.1)

When we use equation (2.8) we get the following tables:

Table 5. Score matrix  $S^1$

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$p_1$	0.7	0.4	0.0	0.6	0.7
$p_2$	0.4	0.6	0.3	0.5	-0.1
$p_3$	0.4	-0.1	0.1	0.8	0.5
$p_4$	0.3	0.5	0.4	0.1	0.3
$p_5$	-0.1	0.4	0.7	-0.1	-0.1

Table 6. Score matrix  $S^2$

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$p_1$	0.4	0.5	0.2	0.4	0.4
$p_2$	0.5	0.4	0.3	0.5	0.4
$p_3$	0.6	0.0	0.5	0.2	0.1
$p_4$	0.6	0.4	0.5	0.7	0.3
$p_5$	0.2	0.4	0.2	0.1	0.4

Table 7. Score matrix  $S^3$

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$p_1$	0.4	0.6	0.6	0.5	0.6
$p_2$	0.7	0.2	0.5	0.1	0.4
$p_3$	0.2	0.3	0.4	0.8	0.7
$p_4$	0.3	0.3	0.2	0.3	0.4
$p_5$	0.3	0.2	0.7	0.3	0.3

Table 8. Score matrix  $S^4$ 

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$p_1$	0.8	0.4	0.7	0.5	0.7
$p_2$	0.7	0.6	0.4	0.2	0.3
$p_3$	0.0	0.2	0.3	0.5	0.6
$p_4$	-0.2	0.6	0.3	0.4	0.5
$p_5$	-0.1	0.2	0.3	0.6	0.5

We use the following equation

$$s_{ij}(w) = \sum_{k=1}^4 w_k s_{ij}^{(k)}$$

and the weight vector  $\omega = (0.4, 0.2, 0.1, 0.3)^T$  to get the following results:

$$\begin{aligned} s_{11}(w) &= \sum_{k=1}^4 w_k s_{11}^{(k)} = w_1 s_{11}^{(1)} + w_2 s_{11}^{(2)} + w_3 s_{11}^{(3)} + w_4 s_{11}^{(4)} \\ &= (0.4)(0.7) + (0.2)(0.4) + (0.1)(0.4) + (0.3)(0.8) = 0.64 \end{aligned}$$

$$\begin{aligned} s_{12}(w) &= \sum_{k=1}^4 w_k s_{12}^{(k)} = w_1 s_{12}^{(1)} + w_2 s_{12}^{(2)} + w_3 s_{12}^{(3)} + w_4 s_{12}^{(4)} \\ &= (0.4)(0.4) + (0.2)(0.5) + (0.1)(0.6) + (0.3)(0.4) = 0.44 \end{aligned}$$

$$\begin{aligned} s_{13}(w) &= \sum_{k=1}^4 w_k s_{13}^{(k)} = w_1 s_{13}^{(1)} + w_2 s_{13}^{(2)} + w_3 s_{13}^{(3)} + w_4 s_{13}^{(4)} \\ &= (0.4)(0.0) + (0.2)(0.2) + (0.1)(0.6) + (0.3)(0.7) = 0.31 \end{aligned}$$

$$\begin{aligned} s_{14}(w) &= \sum_{k=1}^4 w_k s_{14}^{(k)} = w_1 s_{14}^{(1)} + w_2 s_{14}^{(2)} + w_3 s_{14}^{(3)} + w_4 s_{14}^{(4)} \\ &= (0.4)(0.6) + (0.2)(0.4) + (0.1)(0.5) + (0.3)(0.5) = 0.52 \end{aligned}$$

$$\begin{aligned} s_{15}(w) &= \sum_{k=1}^4 w_k s_{15}^{(k)} = w_1 s_{15}^{(1)} + w_2 s_{15}^{(2)} + w_3 s_{15}^{(3)} + w_4 s_{15}^{(4)} \\ &= (0.4)(0.7) + (0.2)(0.4) + (0.1)(0.6) + (0.3)(0.7) = 0.63 \end{aligned}$$

$$s_{21}(w) = \sum_{k=1}^4 w_k s_{21}^{(k)} = w_1 s_{21}^{(1)} + w_2 s_{21}^{(2)} + w_3 s_{21}^{(3)} + w_4 s_{21}^{(4)}$$

$$= (0.4)(0.4) + (0.2)(0.5) + (0.1)(0.7) + (0.3)(0.7) = 0.54$$

$$s_{22}(w) = \sum_{k=1}^4 w_k s_{22}^{(k)} = w_1 s_{22}^{(1)} + w_2 s_{22}^{(2)} + w_3 s_{22}^{(3)} + w_4 s_{22}^{(4)}$$

$$= (0.4)(0.6) + (0.2)(0.4) + (0.1)(0.2) + (0.3)(0.6) = 0.52$$

$$s_{23}(w) = (0.4)(0.3) + (0.2)(0.3) + (0.1)(0.5) + (0.3)(0.4) = 0.35$$

$$s_{24}(w) = (0.4)(0.5) + (0.2)(0.5) + (0.1)(0.1) + (0.3)(0.2) = 0.37$$

$$s_{25}(w) = (0.4)(-0.1) + (0.2)(0.4) + (0.1)(0.4) + (0.3)(0.3) = 0.17$$

We show the values of the first and second rows, in the following table the remaining results can be obtained similarly.

Table 9. Collective score matrix  $S$

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$p_1$	0.64	0.44	0.31	0.52	0.63
$p_2$	0.54	0.52	0.35	0.37	0.17
$p_3$	0.3	0.05	0.27	0.59	0.47
$p_4$	0.21	0.49	0.37	0.33	0.37
$p_5$	0	0.32	0.48	0.19	0.22

Now we use the weight row  $z = (0.2, 0.15, 0.2, 0.3, 0.15)^T$  and the weight function of the score

$$s_j(z_i) = \sum_{i=1}^5 z_i s_{ij} \quad j = 1, 2, 3, 4, 5$$

to obtain the following results:

$$s_1(z_i) = \sum_{i=1}^5 z_i s_{i1} = z_1 s_{11} + z_2 s_{21} + z_3 s_{31} + z_4 s_{41} + z_5 s_{51}$$

$$= (0.2)(0.64) + (0.15)(0.54) + (0.2)(0.3) + (0.3)(0.21) + (0.15)(0.0) = 0.3320$$

$$s_2(z_i) = z_1 s_{12} + z_2 s_{22} + z_3 s_{32} + z_4 s_{42} + z_5 s_{52}$$

$$= (0.2)(0.44) + (0.15)(0.52) + (0.2)(0.05) + (0.3)(0.49) + (0.15)(0.32) = 0.3320$$

$$s_3(z_i) = z_1s_{13} + z_2s_{23} + z_3s_{33} + z_4s_{43} + z_5s_{53}$$

$$= (0.2)(0.31) + (0.15)(0.35) + (0.2)(0.27) + (0.3)(0.37) + (0.15)(0.48) = 0.3515$$

$$s_4(z_i) = (0.2)(0.52) + (0.15)(0.37) + (0.2)(0.59) + (0.3)(0.33) + (0.15)(0.19) = 0.405$$

$$s_5(z_i) = (0.2)(0.63) + (0.15)(0.17) + (0.2)(0.47) + (0.3)(0.37) + (0.15)(0.22) = 0.3895$$

The score function of  $M_j$  is then shown in the following table:

Table 10. The score function of  $M_j$

	$s_1$	$s_2$	$s_3$	$s_4$	$s_5$
$M_j$	0.3320	0.3710	0.3515	0.4050	0.3895

So we can arrange the W.D.F.S.H. as follows:

$$M_1 \leq M_3 \leq M_2 \leq M_5 \leq M_4 \tag{3.14}$$

Equation 3.14 shows that  $M_4$  is the best global supplier because its intuitionistic fuzzy evaluations, after expert-weighted aggregation, give it the highest score on the most heavily weighted criteria—particularly supplier profile and service performance—resulting in the highest overall D.F.S.H. score among all suppliers. So, the best global supplier is  $M_4$ .

All the results in the previous application is confirmed by python coding. The following implementation on decision-making in medical diagnosis.

**Example 2** [53] Let  $U = \{t_1, t_2, t_3, t_4, t_5\}$  be the universal set with D.F.S.H.  $\Theta_1 = \{l_1, l_2, l_3, l_4, l_5\}$  and  $\Theta_2 = \{M_1, M_2, M_3, M_4\}$  where  $l_1, l_2, l_3, l_4, l_5$  are five diseases, such as  $l_1$ (Malaria),  $l_2$ (Typhoid),  $l_3$ (Chestproblem),  $l_4$ (Stomachproblem),  $l_5$ (fever) and  $M_1, M_2, M_3, M_4$  are four patients, the universal set  $U$  represents five symptoms  $t_1$ (Headache),  $t_2$ (Cough),  $t_3$ (Chestpain),  $t_4$ (stomachachpain),  $t_5$ (Bodytempreture), in the following the distance measure  $\delta(\Theta_1, \Theta_2)$  between  $\Theta_1$  and  $\Theta_2$  will be calculated according to our novel distance measure. Decision making is taken according to the fact that the smaller the distance measure, the more likely the patient will be diagnosed with the disease. In the following tables,

Table 11. Disease data for symptoms

	$t_1$	$t_2$	$t_3$	$t_4$	$t_5$
$l_1$	{{0.6, 0.4, 0.3},{0.2, 0.0}}	{{0.7, 0.5, 0.3, 0.2},{0.3, 0.1}}	{{0.5, 0.3},{0.5, 0.4, 0.2}}	{{0.5, 0.4, 0.3, 0.2, 0.1},{0.5, 0.3}}	{{0.5, 0.4, 0.2, 0.1},{0.5, 0.4, 0.3}}
$l_2$	{{0.9, 0.8, 0.7},{0.1, 0.0}}	{{0.5, 0.3, 0.2, 0.1},{0.4, 0.3}}	{{0.2, 0.1},{0.7, 0.6, 0.5}}	{{0.6, 0.5, 0.3, 0.2, 0.1},{0.3, 0.2}}	{{0.4, 0.3, 0.2, 0.1},{0.6, 0.5, 0.4}}
$l_3$	{{0.6, 0.3, 0.1},{0.3, 0.2}}	{{0.9, 0.8, 0.7, 0.6},{0.1, 0.0}}	{{0.5, 0.3},{0.5, 0.4, 0.3}}	{{0.5, 0.4, 0.3, 0.2, 0.1},{0.5, 0.4}}	{{0.6, 0.4, 0.3, 0.2},{0.4, 0.3, 0.2}}
$l_4$	{{0.5, 0.4, 0.2},{0.5, 0.3}}	{{0.4, 0.3, 0.2, 0.1},{0.4, 0.3}}	{{0.4, 0.3},{0.6, 0.5, 0.4}}	{{0.9, 0.8, 0.7, 0.6, 0.5},{0.1, 0.0}}	{{0.5, 0.4, 0.2, 0.1},{0.5, 0.4, 0.3}}
$l_5$	{{0.3, 0.2, 0.1},{0.7, 0.6}}	{{0.5, 0.3, 0.2, 0.1},{0.5, 0.3}}	{{0.3, 0.2},{0.6, 0.4, 0.3}}	{{0.7, 0.6, 0.5, 0.3, 0.2},{0.2, 0.1}}	{{0.8, 0.7, 0.6, 0.5},{0.2, 0.1, 0.0}}

Table 12. Patient data for symptoms

	$t_1$	$t_2$	$t_3$	$t_4$	$t_5$
$M_1$	{{0.9, 0.7, 0.5},{0.1, 0.0}}	{{0.4, 0.3, 0.2, 0.1},{0.5, 0.4}}	{{0.4, 0.3},{0.5, 0.4, 0.2}}	{{0.6, 0.5, 0.4, 0.2, 0.1},{0.3, 0.2}}	{{0.4, 0.3, 0.2, 0.1},{0.5, 0.4, 0.3}}
$M_2$	{{0.5, 0.4, 0.2},{0.5, 0.3}}	{{0.5, 0.4, 0.3, 0.1},{0.4, 0.3}}	{{0.2, 0.1},{0.7, 0.6, 0.5}}	{{0.9, 0.8, 0.6, 0.5, 0.4},{0.1, 0.0}}	{{0.5, 0.4, 0.3, 0.2},{0.5, 0.4, 0.3}}
$M_3$	{{0.9, 0.7, 0.6},{0.1, 0.0}}	{{0.7, 0.4, 0.3, 0.1},{0.2, 0.1}}	{{0.3, 0.2},{0.5, 0.4, 0.3}}	{{0.6, 0.4, 0.3, 0.2, 0.1},{0.4, 0.3}}	{{0.6, 0.3, 0.2, 0.1},{0.4, 0.3, 0.2}}
$M_4$	{{0.8, 0.7, 0.5},{0.2, 0.1}}	{{0.6, 0.5, 0.4, 0.2},{0.4, 0.3}}	{{0.5, 0.3},{0.5, 0.4, 0.3}}	{{0.6, 0.4, 0.3, 0.2, 0.1},{0.4, 0.3}}	{{0.5, 0.4, 0.2, 0.1},{0.5, 0.4, 0.3}}

We used the Python program to compute the following metric when  $\lambda = 0.5$ ,  $\lambda = 1$ ,  $\lambda = 2$ .

Table 13.  $M_1$  when  $\lambda = 0.5$ ,  $\lambda = 1$ ,  $\lambda = 2$ 

$M_1$	$\lambda = 0.5$	$\lambda = 1$	$\lambda = 2$
$l_1$	0.03601762	0.07149409	0.11876774
$l_2$	<b>0.02192978</b>	<b>0.05430682</b>	<b>0.09506508</b>
$l_3$	0.08886402	0.15362159	0.24865306
$l_4$	0.05314720	0.11811907	0.21174163
$l_5$	0.08956279	0.15965574	0.26284623

Table 14.  $M_2$  when  $\lambda = 0.5$ ,  $\lambda = 1$ ,  $\lambda = 2$ 

$M_2$	$\lambda = 0.5$	$\lambda = 1$	$\lambda = 2$
$l_1$	0.08568493	0.14428164	0.22172978
$l_2$	0.04628080	0.11535905	0.21511797
$l_3$	0.10587763	0.16791103	0.25564072
$l_4$	<b>0.01012882</b>	<b>0.03387334</b>	<b>0.06892046</b>
$l_5$	0.07410045	0.12175801	0.18402569

Table 15.  $M_3$  when  $\lambda = 0.5$ ,  $\lambda = 1$ ,  $\lambda = 2$ 

$M_3$	$\lambda = 0.5$	$\lambda = 1$	$\lambda = 2$
$l_1$	<b>0.03758830</b>	<b>0.06440703</b>	<b>0.09801902</b>
$l_2$	<b>0.03693951</b>	0.06977840	0.10415730
$l_3$	0.05813835	0.12072951	0.21444210
$l_4$	0.08973059	0.15219970	0.23604621
$l_5$	0.09387574	0.16814634	0.26850910

Table 16.  $M_4$  when  $\lambda = 0.5, \lambda = 1, \lambda = 2$

$M_4$	$\lambda = 0.5$	$\lambda = 1$	$\lambda = 2$
$l_1$	<b>0.02326248</b>	<b>0.05903791</b>	<b>0.12267776</b>
$l_2$	0.04120925	0.07724478	0.13184612
$l_3$	0.05035189	0.10380775	0.18599472
$l_4$	0.04759524	0.10663432	0.18964704
$l_5$	0.09759155	0.15477417	0.22570673

Based on the above results, we can diagnose  $M_1$  with Typhoid  $l_2$ ,  $M_2$  with the Stomach problem  $l_4$ , and  $M_3, M_4$  with Malaria  $l_1$  as the following table shows:

Table 17. The diagnose by the distance measure

	diagnose
$M_1$	Typhoid $l_2$
$M_2$	Stomach problem $l_4$
$M_3$	Malaria $l_1$
$M_4$	Malaria $l_1$

Now we compare our distance measure with the previous distance measure in [53] when  $\lambda = 1$ : Table 12 shows the values of the old distance measure [53] for  $\lambda = 1$

Table 18. The values of the old distance measure [53]

	$M_1$	$M_2$	$M_3$	$M_4$
$l_1$	0.0712167	0.121351	0.0586095	0.040090
$l_2$	0.050783	0.092528	0.0667410	0.067123
$l_3$	0.127008	0.144407	0.090505	0.0862206
$l_4$	0.0965736	0.030229	0.128719	0.0944345
$l_5$	0.1401883	0.1009896	0.148002	0.141718

From the comparison graphs, it can be observed that the values produced by the proposed distance measure (3.7) are generally slightly higher than those obtained using the classical distance measure[53]. This behavior is expected and can be explained by the structural enrichment of the new distance. Unlike classical distances, which mainly reflect differences in the central tendencies of dual hesitant fuzzy elements, the proposed distance incorporates additional informative components, including the mean function  $\phi$ , the standard deviation  $b$ , the relative coefficient of variation  $c$ , and the hesitancy degree  $\eta$ . As a result, the new distance responds not only to differences in membership and non-membership degrees, but also to discrepancies in dispersion, stability, and hesitation between dual hesitant fuzzy sets. The graphical results indicate that when two alternatives exhibit similar average assessments but differ in their internal variability or degree of hesitation, the proposed distance becomes more sensitive to these differences. This enhanced sensitivity leads to slightly larger distance values in

comparison with the classical measure, thereby providing a more informative and realistic evaluation of dissimilarity. Importantly, the overall ranking patterns in the graphs remain consistent with those obtained by the classical distance, demonstrating that the proposed measure preserves compatibility with existing distance formulations. At the same time, it offers improved discriminatory power in situations characterized by higher uncertainty or greater diversity in expert opinions. Therefore, the proposed distance measure achieves a balance between consistency and sensitivity, extending classical approaches by capturing richer information without causing abrupt changes in decision outcomes. This makes it particularly suitable for complex decision-making problems, such as medical diagnosis, where uncertainty and hesitation play a crucial role.



Figure 2. A comparison between old and new distances for patients

Now we apply the score in (2.4) to diagnose every patient:

Table 19. Score function of disease data for symptoms

	$t_1$	$t_2$	$t_3$	$t_4$	$t_5$
$l_1$	0.333	0.225	0.033	-0.100	-0.100
$l_2$	0.750	-0.075	-0.450	0.090	-0.250
$l_3$	0.083	0.700	0.000	-0.150	-0.075
$l_4$	-0.033	-0.100	-0.150	0.650	-0.100
$l_5$	-0.450	-0.125	-0.183	0.310	0.550

Table 20. Score function of patient data for symptoms

	$t_1$	$t_2$	$t_3$	$t_4$	$t_5$
$M_1$	0.650	-0.200	-0.016	0.110	-0.150
$M_2$	0.033	-0.025	-0.450	0.590	-0.050
$M_3$	0.683	0.225	-0.150	-0.030	0.000
$M_4$	0.517	0.075	0.000	-0.030	-0.100

The following table shows the distance between patients and disease according to the score functions of tables (11,12) where  $D(l_i, M_j) = \sum_{k=1}^5 |\Upsilon(l_i, t_k) - \Upsilon(t_k, M_j)|$  where  $i = 1, 2, 3, 4, 5, j = 1, 2, 3, 4, k = 1, 2, 3, 4, 5$

Table 21. Score function of patient data for symptoms

	$l_1$	$l_2$	$l_3$	$l_4$	$l_5$
$M_1$	1.0517	0.7783	1.9683	1.5067	2.2417
$M_2$	1.8400	1.5333	2.1567	0.4850	1.6633
$M_3$	0.7033	1.0367	1.4200	1.8217	2.4067
$M_4$	0.4367	1.1033	1.3533	1.5550	2.3400

Table 22. The diagnose by the score function

	diagnose
$M_1$	Typhoid $l_2$
$M_2$	Stomach problem $l_4$
$M_3$	Malaria $l_1$
$M_4$	Malaria $l_1$

The results in the above table gives the same result obtained by the distance measure in diagnose, we conclude that the score function is easier in calculations and efficient also in diagnose as well as distance measure We note that the score-function--based solution provides a simplified and efficient decision tool suitable for quick assessments, while the distance-function--based solution offers a richer and more comprehensive evaluation by preserving the intrinsic properties of dual hesitant fuzzy information. The consistency between the two approaches confirms the validity of the proposed distance measure, while the additional discriminatory power of the distance--based method demonstrates its advantage in complex and uncertainty-intensive decision-making environments.

#### 4. Conclusion

From the results presented, we conclude that the proposed distance measure, indicated by  $\delta$ , produces results that are larger than distance in[53], which confirms that the new distance measure clarifies the

data variance. This reflects the importance of the relative coefficient of variation function in calculating the distance function. We find that the new and old distances sometimes are equal, or the new function is larger, which means that the new function reveals data that are more dispersed as well as homogeneous data. Moreover, the auxiliary functions introduced provide valuable interpretive tools: the upper and lower bound functions capture the range of values of W.D.F.S.H., the mean function reflects the average opinion of decision makers, the standard deviation quantifies the degree of dispersion among their opinions, the relative coefficient of variation characterizes the relative variability by relating the mean to the standard deviation, and the hesitancy degree function represents the extent of indecision in the decision-making process. also when compared the solution by score function and distance function, we found that the score-function--based solution provides a simplified and efficient decision tool suitable for quick assessments, while the distance-function--based solution offers a richer and more comprehensive evaluation by preserving the intrinsic properties of dual hesitant fuzzy information. The consistency between the two approaches confirms the validity of the proposed distance measure, while the additional discriminatory power of the distance--based method demonstrates its advantage in complex and uncertainty-intensive decision-making environments We guide the reader to interesting points in [4,5,6,9,22,25].

### Declarations

**Data availability:**Data sets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

**Conflict of Interest:** The authors declare that they have no known competing financial interests or personal relationships that could appear to influence the work reported in this article.

**Funding :** There is no applicable fund

**Authors' contributions :** Kamal Abdelraouf Eldahsha: Writing-original draft, Validation, Software, Methodology, Investigation, Conceptualization. Ghada Mostafa: Writing-original draft, Visualization, Validation, Methodology, Investigation, Conceptualization. Salah Boulaaras and Ibrahim Mekawy: Writing and review & editing, Visualization, Validation, Supervision, Methodology, Investigation, Conceptualization. Zeineb Mohammed: Writing and review & editing, Validation, Methodology, Investigation, Funding acquisition.

### References

1. S.-M. Chen, Measures of similarity between vague sets, *Fuzzy Sets and Systems* 74 (2), 217–223,(1995).
2. S.-M. Chen, J.-M. Tan, Handling multicriteria fuzzy decision-making problems based on vague set theory, *Fuzzy Sets and Systems* 67 163–172, (2) (1994).
3. S.K. De, R. Biswas, A.R. Roy, An application of intuitionistic fuzzy sets in medical diagnosis, *Fuzzy Sets and Systems* 117 209–213,(2) (2001).
4. El-Fawal, Ragaa; Badawy, Abd El-Mohsen; and Hassanein, Abd El-Rahman  $j$ -filters and  $j$ -congruences of locally bounded  $K$  2-algebras, *Al-Azhar Bulletin of Science: Vol. 35: Iss. 3, Article 6,(2024)*.
5. S. A. El-Zahaby and Samira El-Tamimy, Second order optimality conditions for a semilinear elliptic control problem of infinite order with pointwise control constraints, *Journal of Mathematics and Statistical Science (ISSN24112518, USA) vol 5,263-273(2019)*.
6. S. A. El-Zahaby and Samira El-Tamimy, Second order sufficient optimality conditions for some state-constrained control problems of semilinear elliptic equations of infinite order, *Journal of Applied Mathematics And Information Sciences. 18, No. 5, 1075-1080 (2024)*

7. Emanuel O, Patricia M, Oscar C. Comparative study of interval Type-2 and general Type-2 fuzzy systems in medical diagnosis. *Inf Sci.*525:37-53,( 2020)
8. Emanuel OR, Oscar C, Patricia M. Towards asymmetric uncertainty modeling in designing general Type-2 fuzzy classifiers for medical diagnosis. *Expert Syst Appl.* 183:115370-1-12,(2021).
9. Esmaeil, Gehan A. M.; Zaghrout, Afaf A. S.; and El-Deeb, Ahmed A. "Dynamic Ulam stability behavior for first-order nonlinear equations on time scales," *Al-Azhar Bulletin of Science: Vol. 36: Iss. 1, Article 5*,(2025).
10. Farhadinia, B. Information measures for hesitant fuzzy sets and interval-valued hesitant fuzzy sets. *Inf. Sci.* 240, 129–144 (2013).
11. Ghada E. Mostafa, Salah Boulaaras, Rashid Jan, Novel picture fuzzy distance measure  $\rho^*$  and its application in medical diagnosis, *Partial differential equations in applied mathematics*, V.10, 100679, June (2024).
12. Z.-W. Gong, L.-S. Li, F.-X. Zhou, T.-X. Yao, Goal programming approaches to obtain the priority vectors from the intuitionistic fuzzy preference relations, *Computers and Industrial Engineering* 57 1187–1193,(4) (2009).
13. M. Grabisch, S. Greco, M. Pirlot, Bipolar and bivariate models in multicriteria decision analysis: descriptive and constructive approaches, *International Journal of Intelligent Systems* 23 930–969,(9) (2008).
14. D.H. Hong, C. Kim, A note on similarity measures between vague sets and between elements, *Information Sciences* 115 (1–4) (1999) 83–96.
15. A. Kharal, Homeopathic drug selection using intuitionistic fuzzy sets, *Homeopathy* 98 35–39,(1) (2009).
16. Hoseini J, Mohammadi A Estimator and Tests for Coefficient of Variation in Uniform Distribution. *J Biomet Biostat* 3:149.(2012).
17. D.-F. Li, G.-H. Chen, Z.-G. Huang, Linear programming method for multiattribute group decision making using IF sets, *Information Sciences* 180 1591–1609, (9) (2010).
18. Li, D. Q., Zeng, W. Y. , Li, J. H. New distance and similarity measures on hesitant fuzzy sets and their applications in multiple criteria decision making. *Eng. Appl. Artif. Intell.* 40, 11–16 (2015).
19. Li, D. Q., Zeng, W. Y. , Zhao, Y. B. Note on distance measure of hesitant fuzzy sets. *Inf. Sci.* 321, 103–115 (2015).
20. L. Lin, X.-H. Yuan, Z.-Q. Xia, Multicriteria fuzzy decision making methods based on intuitionistic fuzzy sets, *Journal of Computer and System Sciences* 73 84–88, (1) (2007).
21. H.-W. Liu, G.-J. Wang, Multi-criteria decision-making methods based on intuitionistic fuzzy sets, *European Journal of Operational Research* 179 (1) 220–233, (2007).
22. Mahmoud, Asmaa M.; Eid, Heba F.; Desuky, Abeer S.; and Ali, Hoda A. "Graph neural network guided by feature selection and centrality measures for Node Classification on Homophilic and Heterophily graphs," *Al-Azhar Bulletin of Science: Vol. 35: Iss. 1, Article 2*, (2024).
23. Molla MU, Giri BC, Biswas P. Extended PROMETHEE method with Pythagorean fuzzy sets for medical diagnosis problems. *Soft Comput.* 25:4503-4512,(2021).
24. Boulaaras, S., Mostafa, G.E., Jan, R. et al. Applications of distance measure between dual hesitant fuzzy sets in medical diagnosis and weighted dual hesitant fuzzy sets in making decision. *Sci Rep* 14, 26214 (2024).
25. Samira El-Tamimy, Linear elliptic control problem of infinite order with pointwise state constraint, *Journal of Applied Mathematics And Information Sciences.* 16, No 6, 1005-1008 (2022).
26. Singh P. Distance and similarity measures for multiple-attribute decision making with dual hesitant fuzzy sets. *J Comput Appl Math.*36:1-16,(2015)
27. Su Z, Xu ZS, Liu HF, Liu SS. Distance and similarity measures for dual hesitant fuzzy sets and their applications in pattern recognition. *J Intell Fuzzy Syst.* ;29:731-745,(2015).

28. Tamalika Chaira, A new measure using intuitionistic fuzzy set theory and its application to edge detection, *Applied soft computing*, V.8,p.919-927,(2008).
29. C. Tan, X. Chen, Intuitionistic fuzzy Coquet integral operator for multi-criteria decision making, *Expert Systems with Applications* 37 149-157, (1) (2010).
30. Ting-Yu Chen, A comparative analysis of score functions for multiple criteria, decision making in intuitionistic fuzzy settings, *Information Sciences*, Volume 181, Issue 17, Pages 3652-367,1 September (2011).
31. Torra V. Hesitant fuzzy sets. *Int J Intell Syst.* 25:529-539,(2010).
32. Torra V, Narukawa Y. On hesitant fuzzy sets and decision. In: Hong TJ, Kyung CM, Kyung WO, eds. *The 18th IEEE International Conference on Fuzzy Systems*, Jeju Island, Korea, IEEE; 1378-1382,(2009).
33. A. Tversky, D. Kahneman, *Advances in prospect theory: cumulative representation of uncertainty*, *Journal of Risk and Uncertainty* 5 297-323,(4) (1992).
34. Wang RG, Li WM, Zhang T, Han Q. New distance measures for dual hesitant fuzzy sets and their application to multiple attribute decision making. *Symmetry-Basel* 12:191-208,(2020).
35. P. Wang, QoS-aware web services selection with intuitionistic fuzzy set under consumer's vague perception, *Expert Systems with Application* 36 4460–4466, (3) (2009).
36. G. Wei, Some induced geometric aggregation operators with intuitionistic fuzzy information and their application to group decision making, *Applied Soft Computing* 10 423–431, (2) (2010).
37. G.-W. Wei, GRA method for multiple attribute decision making with incomplete weight information in intuitionistic fuzzy setting, *Knowledge-Based Systems* 23 243–247, (3) (2010).
38. Xia MM, Xu ZS. Hesitant fuzzy information aggregation in decision making. *Int J Approx Reasoning.*52:395-407,(2011).
39. Xia MM, Xu ZS, Chen N. Some hesitant fuzzy aggregation operators with their application in group decision making. *Group Decis Negotiation.* 22:259-279,(2013).
40. Xu ZS, Xia MM. Hesitant fuzzy entropy and cross-entropy and their use in multiattribute decision-making. *Int J Intell Syst.* 27:799-822,(2012).
41. Xu ZS, Xia MM. Distance and similarity measures for hesitant fuzzy sets. *Inf Sci.* 181:2128-2138,(2011).
42. Xu ZS, Xia MM. On distance and correlation measures of hesitant fuzzy information. *Int J Intell Syst.*26:410-425,(2011).
43. Z. Xu, Methods for aggregating interval-valued intuitionistic fuzzy information and their application to decision making, *Control and Decision* 22 215–219, (2) (2007).
44. Z. Xu, Intuitionistic preference relations and their application in group decision making, *Information Sciences* 177 2363–2379, (11) (2007).
45. Z. Xu, Intuitionistic fuzzy aggregation operators, *IEEE Transactions on Fuzzy Systems* 15 1179–1187, (6) (2007).
46. Z. Xu, Multi-person multi-attribute decision making models under intuitionistic fuzzy environment, *Fuzzy Optimization and Decision Making* 6 221–236,(3)(2007).
47. Z. Xu, Choquet integrals of weighted intuitionistic fuzzy information, *Information Sciences* 180 726–736,(5) (2010).
48. Z.-S. Xu, J. Chen, Approach to group decision making based on interval-valued intuitionistic judgment matrices, *Systems Engineering-Theory, Practice* 27 126–133, (4) (2007).
49. Z. Xu, R.R. Yager, Some geometric aggregation operators based on intuitionistic fuzzy sets, *International Journal of General Systems* 35 417–433, (4) (2006).
50. Z.S. X.u, Multi-person multi-attribute decision making models under intuitionistic fuzzy environment, *Fuzzy Optim. Decis.* 221-236, Making 6(2007).
51. Zadeh LA. Fuzzy sets. *Inf Control.* 8:338-356,(1965).
52. L.A. Zadeh Concept of a linguistic variable and its application to approximate reasoning, I, II, III *Informat. Sci.*, 8, pp. 199–249, 301–357, 43–80(1985).

53. Zeng W, Ma R, Liu Z, Xi Y, Yin Q, Xu Z. Som novel distance measures between dual hesitant fuzzy sets and their application in medical diagnosis. *Int J Intel syst.*1-19(2022).
54. Zeng WY, Xi Y, Yin Q, Guo P. Weighted dual hesitant fuzzy set and its application in group decision making. *Neurocomputing.*458:714-726,(2021)
55. Zeng WY, Li DQ, Yin Q. Distance and similarity measures of hesitant fuzzy sets and their application in pattern recognition. *Pattern Recognit Lett.* 84,267-271,(2016)
56. H. Zhao, Z. Xu, M. Ni, S. Liu, Generalized aggregation operators for intuitionistic fuzzy sets, *International Journal of Intelligent Systems* 25 1–30, (1) (2010).